

8

Urban Ecosystems and Biodiversity

Coordinating Lead Authors

Timon McPhearson (New York), Madhav Karki (Kathmandu)

Lead Authors

Cecilia Herzog (Rio de Janeiro), Helen Santiago Fink (Vienna/D.C.), Luc Abbadie (Paris), Peleg Kremer (New York), Christopher M. Clark (Washington D.C.), Matthew I. Palmer (New York), Katia Perini (Genoa)

Contributing Authors

Marielle Dubbeling (Leusden)

This chapter should be cited as

McPhearson, T., Karki, M., Herzog, C., Santiago Fink, H., Abbadie, L., Kremer, P., Clark, C. M., Palmer, M. I., and Perini, K. (2018). Urban ecosystems and biodiversity. In Rosenzweig, C., W. Solecki, P. Romero-Lankao, S. Mehrotra, S. Dhakal, and S. Ali Ibrahim (eds.), *Climate Change and Cities: Second Assessment Report of the Urban Climate Change Research Network*. Cambridge University Press. New York. 257–318

Urban Ecology in a Changing Climate

Almost all of the impacts of climate change have direct or indirect consequences for urban ecosystems, biodiversity, and the critical ecosystem services they provide for human health and well-being in cities. These impacts are already occurring in urban ecosystems and their constituent living organisms.

Urban ecosystems and biodiversity have an important and expanding role in helping cities adapt to and mitigate the impacts of changing climate. Harnessing urban biodiversity and ecosystems as adaptation and mitigation solutions will help achieve more resilient, sustainable, and livable outcomes for cities and urban regions.

Conserving, restoring, and expanding urban ecosystems under mounting climatic and non-climatic urban development pressures will require improved urban and regional planning, policy, governance, and multisectoral cooperation.

Major Findings

- Urban biodiversity and ecosystems are already being affected by climate change.
- Urban ecosystems are rich in biodiversity and provide critical natural capital for climate change adaptation and mitigation.
- Climate change and urbanization are likely to increase the vulnerability of biodiversity hotspots, urban species, and critical ecosystem services.

- Urban ecosystems and green infrastructure can provide cost-effective, nature-based solutions for adapting to climate change while also creating opportunities to increase social equity, green economies, and sustainable urban development.
- Investing in the quality and quantity of urban ecosystems and green infrastructure has multiple co-benefits, including improving quality of life, human health, and social well-being.

Key Messages

Cities should take a long-term, system-based approach to climate adaptation and mitigation. Nature-based approaches to address climate change in cities explicitly recognize the critical role of urban and peri-urban ecosystem services (UES) that require thoughtful management in order to ensure sustainable supply of environmental goods and services to residents who need them over the next 20, 50, and 100 years. Ecosystem-based planning can strengthen the linkages between urban, peri-urban, and rural ecosystems through participatory planning and management for nature-based solutions at both city and regional scales.

The economic benefits of urban biodiversity and ecosystem services should be quantified so that they can be integrated into climate-related urban resilience and sustainability planning and decision-making. These benefits should incorporate both monetary and non-monetary values of biodiversity and ecosystem services, including how they relate to physical and mental health and social equity in access to services.

8.1 Introduction

Climate change is already affecting cities and urbanized regions around the world impacting human populations and the built environment, as well as urban ecosystems and their associated biota (While and Whitehead 2013). Almost all of the impacts of climate change have direct or indirect consequences for urban ecosystems,¹ biodiversity,² and the critical ecosystem services³ they provide for human health and well-being in cities (e.g., urban heat island [UHI] reduction) (The Economics of Environmentalism and Biodiversity [TEEB], 2011; Elmqvist et al., 2013). Increasing knowledge of the benefits of urban ecosystems for the livelihoods of urban residents suggests an important and expanding role for urban ecosystems and biodiversity in adaptation to local effects of climate change. However, conserving, restoring, and expanding urban ecosystems to enhance climate resilience and other co-benefits under mounting climatic and non-climatic stresses of growing urbanization and development processes will require improved urban and regional planning, policy and governance, and multisectoral cooperation to protect and manage urban ecosystems and biodiversity (Elmqvist et al., 2013; Solecki and Marcotullio, 2013; McPhearson et al., 2014).

In this chapter, we review key concepts, challenges, and ecosystem-based pathways for adaptation and mitigation of climate change in cities. This leads to and supports concepts, strategies, and tools of ecosystem-based adaptation, disaster risk reduction, and green infrastructure planning. Section 8.1 reviews the relationships among urban ecosystems, biodiversity, and ecosystem services as critical resources for climate adaptation and, to some extent, mitigation. Sections 8.2 and 8.3 discuss current and future climate-related challenges including hazards, risks, and vulnerabilities for urban biodiversity and ecosystem services. Section 8.4 discusses examples of how ecosystems can provide adaptive capacity and be used innovatively to reduce effects of climate change in urban systems, whereas Section 8.5 presents ecosystem-based adaptation as an effective entry point for nature-based solutions to building climate resilience in cities. Section 8.6 discusses the economic cost-effectiveness of ecosystem-based adaptation, with particular emphasis on investing in green infrastructure. Section 8.7 discusses how urban ecosystems intersect with urban planning and design (see Chapter 5, Urban Planning and Design), the importance of engaging with diverse stakeholders, and how ecosystem-based planning and management can help address issues of social equity and environmental justice while yielding multiple socioeconomic benefits. Section 8.8 discusses important planning, governance, and management tools (see Chapter

16, Governance and Policy). Sections 8.9 and 8.10 present the need for better linking science with policy, in particular for building urban climate resilience. Section 8.11 identifies remaining knowledge gaps and suggests avenues for future research. Section 8.12 provides a summary of recommendations for cities to harness urban biodiversity and ecosystems as nature-based solutions to adapt to the effects of and mitigate climate change that will help achieve more sustainable, resilient, and livable cities. Case Studies are provided throughout the chapter to illustrate effective, on-the-ground implementation of many of the ecosystem-based adaptation and mitigation strategies and approaches reviewed.

8.1.1 A Systems Approach to Ecology in, of, and for Cities

Cities and urban areas are complex systems with social, ecological, economic, and technical/built components interacting dynamically in space and time (Grimm et al., 2000, 2008; Pickett et al., 2001; McPhearson et al., 2016a). The complex nature of urban systems⁴ can make it challenging to predict how ecosystems will respond to climate change in cities (Batty 2008; Bettencourt and West, 2010; McPhearson et al., 2016b). This complexity is driven by many intersecting feedbacks affecting ecosystems, including climate, biogeochemistry, nutrient cycling, hydrology, population growth, urbanization and development, human perceptions and behavior, and more (Bardsley and Hugo, 2010; Pandey and Bardsley, 2013; Alberti, 2015).

In urban ecology, cities and urbanized areas are understood to be complex human-dominated ecosystems (Pickett et al., 1997, 2001; Niemelä et al., 2011). These systems interrelate dynamically with the social, ecological, economic, and technological/built infrastructure of the city (Grimm et al., 2000; McDonnell and Hahs, 2013; McPhearson et al., 2016a) (see Figure 8.1). Patterns and processes of urban systems in this view emerge from the interactions and feedbacks between components and systems in cities, emphasizing the need to consider multiple sources of social-ecological patterns and processes to understand reciprocal interactions between climate change and urban ecosystems (see Figure 8.1).

Urban social-ecological systems (SES) consist of social and ecological components (broadly defined) that have their own internal patterns and processes, but these patterns and processes interact across the system in a number of ways to produce overall urban system dynamics, behavior, and emergent phenomena. Drivers external to the urban system are fundamentally important but can affect social and ecological components and processes within the urban system with different strengths

1 Urban ecosystems include all vegetation, soil, and water-covered areas that may be found in urban and peri-urban areas at multiple spatial scales (parcel, neighborhood, municipal city, metropolitan region), including parks, cemeteries, lawns and gardens, green roofs, urban allotments, urban forests, single trees, bare soil, abandoned or vacant land, agricultural land, wetlands, streams, rivers, lakes, and ponds (Gómez-Baggethun et al., 2013).

2 "Biological diversity" means the variability among living organisms from all sources including, inter alia, terrestrial, marine, and other aquatic ecosystems and the ecological complexes of which they are part; this includes diversity within species, between species, and of ecosystems.

3 Ecosystem services are the benefits that people obtain directly or indirectly from ecosystem functions, such as protection from storm surges and heat waves, air quality regulation, and food, fiber, and fresh water (MA, 2005; TEEB, 2010; Gomez-Baggethun et al., 2013).

4 Urban systems are defined here as those areas where the built infrastructure covers a large proportion of the land surface or those in which people live at high densities (Pickett et al., 2001).

URBAN SOCIAL- ECOLOGICAL SYSTEM




Figure 8.1 Urban systems are complex and dynamically interactive and can be conceptualized and studied as social-ecological systems (SES) at multiple spatial or temporal scales.

or intensity. This conceptual approach to studying urban SES is scale-independent and can therefore be applied at multiple spatial or temporal scales in urban areas. The urban ecosystem approach has developed rapidly in the past two decades, incorporating methods and approaches from the social sciences, biophysical sciences, urban planning, and design to provide insight for developing and managing urban areas to meet the needs of expanding populations in a changing climate (McDonnell, 2011; McPhearson et al., 2016a). We focus here on biodiversity and ecosystem functions and services provided by natural systems within urban and peri-urban areas.

Studies of the ecology *in* the city as well as ecology *of* the city (Grimm et al., 2000; Pickett et al., 2001) are both domains of urban ecology, a science increasingly focused on applying sustainability and resilience science *for* cities (Childers et al., 2014, 2015; McPhearson et al., 2016a). Defining clear boundaries for ecosystems *in* the city is challenging due to the fact that species and many of the relevant fluxes and interactions necessary to understand the functioning of urban ecosystems extend beyond the city boundaries defined by political borders (Solecki and Marcotullio, 2013). For example, nutrients, water, species, and humans all move across political boundaries, emphasizing the importance

of regional planning and management. Thus, the relevant scope of urban ecosystem analysis reaches far beyond the municipal boundary. It comprises not only the ecological areas within cities, but also the peri-urban areas and linkages to nearby rural areas that are directly affected by the energy and material flows from the urban core, including city water catchments, peri-urban forests, and nearby cultivated fields (Grimm et al., 2000; Pickett et al., 2001; La Rosa and Privitera, 2013). Urban ecosystems therefore include all vegetation, soil, and water-covered areas that may be found in urban and peri-urban areas at multiple spatial scales (parcel, neighborhood, municipality, metropolitan region), including parks, cemeteries, lawns and gardens, green roofs, urban allotments, urban forests, single trees, bare soil, abandoned or vacant land, agricultural land, wetlands, streams, rivers, lakes, and ponds (Gómez-Baggethun et al., 2013).

The social and biophysical context of urban areas influences resilience to climate change and other social-ecological challenges (Marcotullio and Solecki, 2013; Solecki and Marcotullio, 2013). For example, the bio-geophysical context of the city or urban area may determine how ecosystems in cities respond to climate change, extreme events, and urbanization (Schewenius et al., 2014). Urbanization and suburbanization in urban areas

often reduce both species richness (i.e., the number of species) and evenness (i.e., the distribution of species) for most biotic communities (Paul et al., 2001; McKinney, 2002). Changes in species richness and evenness have been found to affect the stability of ecosystems and their ability to deliver needed services for mitigating and adapting to climate change (Grimm, 2008; Cardinale et al., 2012). Additionally, many of the changes taking place in urban areas have analogues to those driven by climate change (e.g., elevated CO₂, higher temperatures, changes in precipitation), thus making urban systems useful models for examining the interaction of social and biophysical patterns and processes in changing climates (Grimm et al., 2008; Collins et al., 2000). Therefore, urban ecological approaches to improving climate adaptation and mitigation should employ a systems approach characterized by interdisciplinary, multiscale studies and a focus on interactions and feedbacks to further develop an ecology *of* and *for* cities (see Figure 8.1) (Grimm et al., 2000; Pickett et al., 2001; McDonald et al., 2013; Childers et al., 2014, 2015; McPhearson et al., 2016). Green infrastructure and ecosystem-based adaptation are important components of nature-based solutions for climate mitigation and adaptation.

8.1.2 Urban Green Infrastructure

Many cities have already made significant progress employing urban ecological resources for climate change adaptation and mitigation as part of urban infrastructure design, planning, and development (Frischenbruder and Pellegrino, 2006). Green infrastructure is becoming a widely utilized nature-based solution for climate change adaptation and mitigation in cities (Flørgard, 2007). We consider green infrastructure as a network of natural and semi-natural areas, features, and green spaces in rural and urban, terrestrial, coastal, and marine areas, which together enhance ecosystem health and climate change resilience, contribute to biodiversity, and benefit human populations through the maintenance and enhancement of ecosystem services (Pauleit et al., 2011; Kopperoinen et al., 2014). Green infrastructure is often also examined as a specific management tool for combining engineered and ecological systems (e.g., bioswales) in place of engineered non-ecological systems (e.g., concrete sewer drains) to provide ecosystem services such as cooling, stormwater management, UHI reduction, carbon storage, flood protection, and recreation (Novotny et al., 2010).

Urban green infrastructure is a key strategy for mitigating and adapting to the effects of climate change. For example, the UHI effect can be reduced by several degrees through enhanced transpiration and the shading provided by street trees, green roofs, and parks (Onishi et al., 2010; Petralli et al., 2006; Rosenzweig et al., 2006; Susca et al., 2011; Taha, 1997). Vegetation also decreases energy use for heating and air conditioning (McPhearson et al., 1997; Akbari et al., 2001, UNEP, 2011). Akbari et al. (2001) estimated that about 20% of the national cooling demand in the United States can be avoided through a large-scale implementation of heat-island mitigation measures such as urban green infrastructure, particularly urban forestry. Vegetation also contributes to a city's mitigation efforts by capturing CO₂

through photosynthesis and absorbing atmospheric pollutants through dry deposition on leaves and branches, and uptake by stomata (Fowler, 2002; Ottel   et al., 2010). Green roofs and vegetated areas, including trees, increase rainwater infiltration and reduce peak flood discharge and associated water pollution while also providing mental and physical health benefits such as providing spaces for recreation and relaxation and decreasing the level of citizen stress (Dunnett and Kingsbury, 2008; Scholz-Barth, 2001; Czemi  l Berndtsson, 2010; Carson et al., 2013) (see Figure 8.2).

New York, for example, launched the Green Infrastructure Plan in 2010 designed to invest in new and restored green infrastructure for stormwater management instead of traditional gray infrastructure. This included committing US\$1.5 billion for green infrastructure development over the next 20 years (NYC Environmental Protection, 2010) (see Case Study 8.2). Similarly, the city of Taizhou, China, located on the southeast coast of Zhejiang Province with 5.5 million inhabitants, developed a zoning plan that utilized green infrastructure to adapt urban growth to deal with potential impacts of climate change including preventing stormwater related floods and maintaining food production areas (Yu and Li, 2006). The Taizhou plan incorporated ecological areas at multiple scales (local to regional) to maintain critical natural processes and flows including hydrology and biodiversity while simultaneously protecting cultural heritage sites and recreation areas (Yu and Li, 2006; Ahern, 2007; Gotelli et al., 2013). These and other relevant Case Studies described in this chapter demonstrate the importance and cost-effective benefits of incorporating urban ecosystems explicitly into urban design, management, planning, and policy for mitigating and adapting to the effects of climate change.

8.1.3 Urban Biodiversity and Ecosystem Services

Nature in cities plays a crucial role as the ecological basis for human–nature interactions and the production of Urban Ecosystem Services (UES) (see Box 8.1 and Figure 8.3) (Kowarik 2005; Bolund and Hunhammar, 1999; G  mez-Baggethun et al., 2013; TEEB, 2011; Kremer et al., 2016a). Biodiversity and ecosystems in cities are increasingly linked to human health and well-being, livability, and the quality of urban life (McGranahan et al., 2005; G  mez-Baggethun et al., 2013; McPhearson et al., 2013). For example, urban trees can remove harmful air pollution, provide shade during heat waves, absorb and store carbon, and create spaces for contemplation, aesthetic and spiritual enjoyment, and social cohesion (see Table 8.1 and Figure 8.2) (TEEB, 2011; McPhearson 2011; G  mez-Baggethun et al., 2013; Andersson et al., 2014; Nowak et al., 2013).

Biodiversity is the fundamental basis for the generation of ecosystem services (see Figure 8.3) (Elmqvist et al., 2013; G  mez-Baggethun et al., 2013). There are many ecosystem services that cannot be imported and must be supplied locally within urban ecosystems (McPhearson et al., 2013b, 2014; Andersson et al., 2015a). For example, utilizing urban parks, green walls and roofs, and street trees to adapt to and mitigate impacts of

Case Study 8.1 Coastal Natural Protected Areas in Mediterranean Spain: The Ebro Delta and Empordà Wetlands

Sandra Fatorić and Ricard Morén-Alegret

Department of Geography, Autonomous University of Barcelona

Christos Zografos

Institute of Environmental Science and Technology (ICTA), Autonomous University of Barcelona

Keywords	Sea level rise, vulnerability, ecosystems-based adaptation, coastal natural protected area
Population (Study Region)	Ebro Delta: 48,031 Empordà wetlands: 43,354 (IDESCAT, 2015)
Area (Study Region)	Ebro Delta: 299.4 km ² Empordà wetlands: 123 km ² (IDESCAT, 2015)
Income per capita	US\$27,520 (World Bank, 2017)
Climate zone	Csa – Temperate, dry summer, hot summer (Peel et al., 2007)

Climate change is an increasingly significant global problem with potentially far-reaching consequences for coastal human communities, livelihoods, and ecosystems in the Mediterranean region. Seven economically, socially, and environmentally dynamic urban towns across coastal natural protected areas in Mediterranean Spain, the Ebro Delta (see Case Study 8.1 Figure 1), and Empordà wetlands have been particularly vulnerable to three aspects of climate change: (1) air and sea temperature rise (2) sea level rise, and (3) decreased river flows (see Case Study 8.1 Table 1). In addition, intensification of coastal erosion, flooding, saltwater intrusion, and deficits in river sediment supply have been affecting natural habitats and livelihoods in these areas (Barnolas and Llasat, 2007; Candela et al., 2007; CIIRC, 2010; Day et al., 2006; Guillén and Palanques 1992; Jiménez et al., 1997; Martín-Vide et al., 2012; Sánchez-Arcilla et al., 2008).

This Case Study is based on studies that identified a local dimension of climate change adaptation relevant for maintaining a wide range of livelihoods while facing current and future climate change (Fatorić, 2010, 2014). These studies build on the work of Smit and Wandel (2006) who highlighted that adaptation is an outcome of the interaction of environmental, social, cultural, political, and economic forces. Analytically, adaptation is conceptualized in this paper as a set of technical options to respond to specific risks (Nelson et al., 2007), where the need for local stakeholder involvement has been increasingly acknowledged (Bormann et al., 2012; Cote et al., 2014; Eriksen et al., 2011; Mozumder et al., 2011). Different stakeholders may hold different knowledge, opinions, and understandings of the local context of adaptation. Thus, which specific sources of knowledge are recognized and used in decision-making process is crucial for determining which interests, development paths, and solutions are prioritized (Eriksen et al., 2011).

Local, regional, and national stakeholders belonging to various economic sectors (e.g., employees of local tourist information centers,



Case Study 8.1 Figure 1 Locations of the three municipalities in the Ebro Delta and coastal natural protected area.

farmers, peasants, engineers); public administrations (e.g., governmental officers), environmental organizations and research centers (e.g., members of environmental groups, scientists), and social organizations (e.g., members of social and ethnic organizations) linked to the Ebro Delta and Empordà wetlands were selected to participate in the studies.

The results showed that adaptation appears to have taken place in the Ebro Delta and Empordà wetlands during the past few decades, but mainly through unsustainable measures (e.g., artificial or hard structures).

More than half of interviewed stakeholders reported that they favor “natural” adaptation measures such as (1) building and/or restoring coastal sand dunes and (2) raising ground level. Approximately one-quarter were in favor of “artificial” adaptation measures such as (1) seawalls, groins, and breakwaters; (2) flood and underwater gates; (3) beach nourishment; and (4) rainwater harvesting. The remaining stakeholders considered combining both types of measures. Stakeholders were also asked to consider coastal relocation as an adaptation option.

With respect to natural adaptation measures, building sand dunes parallel to the shoreline where none exists and/or restoring and stabilizing the existing ones was perceived as the optimal adaptation measure in both protected coastal areas. This option was often considered as the cheapest one for both study areas, and it is compatible with environmental sustainability actions. Moreover, building and restoring dunes can increase socioecological resilience in both areas and produce benefits in the absence of climate change effects.

Regarding raising ground level, the other natural adaptation measure, interviewees expressed little support for elevating ground level by a few centimeters. This might be due to weak technical and urban design skills among most stakeholders.

Artificial measures, on the other hand, did not have such unified support among stakeholders. For instance, dykes, seawalls, and breakwaters generated different opinions. About one-third of stakeholders were against “artificialization” mainly due to the current ecological value of both areas. Stakeholders perceived these measures as too costly to build and maintain. A small number of stakeholders were willing to preserve an already attractive landscape for economic activities (especially tourism) by implementing artificial measures.

Flood and saltwater intrusion gates were suggested and discussed, but gates may be not suitable measures because they entail significant investments. Beach nourishment was perceived by a minority of stakeholders as a suitable measure that

is aesthetically pleasing and that sometimes can be implemented with a reasonable budget.

The studies also revealed that rainfall capture and storage in those parts of Mediterranean Spain where precipitation is likely to decrease and become more variable has not yet been prioritized.

Regarding coastal relocation, it was interesting to note a difference between the two study areas: according to population data (IDESCAT, 2015), the rate of registered foreign immigrants is higher in Empordà wetlands than in the Ebro Delta, and, interestingly, among the interviewed stakeholders in Empordà wetlands there was more willingness for relocation elsewhere as an “adaptation” measure than in the Ebro Delta. In this sense, it emerged that place

Case Study 8.1 Table 1 Comparison of socioeconomic, environmental, and climatic characteristics of the Ebro Delta and Empordà wetlands. Source: ACA, 2014; BirdLife International, 2014a and 2014b; CREA, 2013; Ebre Observatory, 2015; IDESCAT, 2015; Meteo Estartit, 2015; Ninyerola et al., 2004; SMC, 2015; Wetlands International, 1992 and 1995

	Ebro Delta	Empordà wetlands
<i>Municipalities</i>	Amposta, Deltebre, Sant Carles de la Rapita	Castello d'Empuries, Escala, Roses, Sant Pere Pescador
<i>Physical territory</i>	Coastal lagoons, marshlands, beaches, dunes, salt pans	Coastal lagoons, inland freshwater ponds, marshlands, beaches, dunes
<i>Protection</i>	Ramsar Convention on Wetlands (1986), Natura 2000, Special Protection Area for Birds	Ramsar Convention on Wetlands (1992), Natura 2000, Special Protection Area for Birds
<i>Protected area</i>	11,530 ha	10,830 ha
<i>River(s)</i>	Ebro	Muga, Fluvia
<i>River regulation</i>	Mequinea, Flix, Riba-Roja dams	Boadella dam
<i>Vegetation</i>	<i>Arthrocnemum fruticosi</i> , <i>Crucianellum maritima</i> , <i>Scirpetum maritimi-littoralis</i> , <i>Agropyretum mediterraneum</i>	<i>Salix alba</i> , <i>Fraxinus angustifolia</i> , <i>Rosa sempervivens</i> , <i>Arum italicum</i> , <i>Arista lochia rotunda</i> , <i>Typha latifolia</i>
<i>Fauna</i>	<i>Anas strepera</i> , <i>Phoenicopterus roseus</i> , <i>Botaurus stellaris</i> , <i>Ardea purpurea</i> , <i>Larus audouinii</i>	<i>Coracias garrulus</i> , <i>Lanius minor</i> , <i>Buteo buteo</i> , <i>Falco subbuteo</i> , <i>Bos taurus domestica</i>
<i>Climate</i>	Mediterranean	Mediterranean
<i>Air temperature (1990–2014)</i>	Increase by 0.37°C/decade (Ebre Observatory)	Increase by 0.19°C/decade (Sant Pere Pescador)
<i>Precipitation (1990–2014)</i>	Slight increase (Ebre Observatory)	Decrease (Sant Pere Pescador)
<i>River flow (1990–2012)</i>	Decrease by 14 m ³ /s/decade (Ebro)	Decrease by 3.5 m ³ /s/decade (Fluvia)
<i>Sea level (1990–2014)</i>	Increase by 3.9 cm/decade (Estartit)	Increase by 3.9 cm/decade (Estartit)
<i>Sea temperature (1990–2014)</i>	Increase by 0.18°C (sea surface), 0.17°C (20 m), 0.28°C (50 m), 0.13°C (80 m) per decade (Estartit)	Increase by 0.18°C (sea surface), 0.17°C (20 m), 0.28°C (50 m), 0.13°C (80m) per decade (Estartit)
<i>Tourism</i>	Ecotourism, birdwatching	Campsites, second homes, hotels, ecotourism, birdwatching, marina
<i>Agriculture</i>	Rice, citrus fruits	Fruits trees, vines, olives
<i>Fishery</i>	14% of Catalonia's fish production	11% of Catalonia's fish production
<i>N° population (2014)</i>	48,031	43,354

attachment (and previous migration experience) among local residents is relevant when considering relocation as an adaptation measure.

One of the lessons that can be drawn from this Case Study is the need to gather and integrate local understanding, perception, and knowledge with scientific knowledge in order to develop successful response to climate change, empower local decision-making, and preserve current ecosystems, livelihoods, and communities. Encouraging local communities and policy-makers to undertake short- and medium-term thinking and to develop adaptation planning with more desirable sustainable outcomes should be a priority in the Ebro Delta and Empordà wetlands. Stakeholders can help to

raise awareness in order to implement adaptation measures based on technical solutions that would reduce the vulnerability of natural and socioeconomic systems and take advantage of any potential opportunities and benefits (Fatorić and Chelleri, 2012; Fatorić and Morén-Alegret, 2013; Fatorić et al., 2014).

Another lesson that emerged from the research is that the optimal adaptation measure according to the stakeholders in both coastal protected areas is building and/or restoring coastal dunes, which is likely to be the most efficient and least expensive protection against various climate change effects (see Case Study 8.1 Figure 2). This highlights the need for dune conservation and maintenance as climate change reinforces the value of its protective capacity.



Case Study 8.1 Figure 2 Dunes as “natural” adaptation measure in Empordà wetlands.




Figure 8.2 Urban trees and other types of green infrastructure, shown here from New York, can provide important climate adaptation and mitigation in cities. The quantity and quality of benefits (e.g., carbon storage, urban heat island mitigation, stormwater absorption) depend on the urban context and configuration of trees in an urban landscape. A. Individual scattered street trees at 125th Street and Madison Avenue, Harlem. B. Dense street trees at Eastern Parkway and Classon Avenue, Brooklyn. C. Street trees as a corridor connecting small- and medium-sized urban green spaces, Elmhurst, Juniper Valley Parks, and Lutheran All Faiths Cemetery, Queens. D. Dense urban street trees connecting large urban parks: Bronx Park and Van Cortlandt Park. E. Disconnected urban green space with scattered trees, Green-Wood Cemetery, Brooklyn. F. Large urban forest, Forest Park, Queens.

climate change such as urban heat must occur locally (Gill et al., 2007; Pataki et al., 2011). Urban ecosystems are therefore especially important in delivering climate-related ecosystem services with direct impact on human health, well-being, and security (Novotny et al., 2010; Elmqvist et al., 2013; McPhearson et al., 2015). Additionally, investing in urban ecosystems for climate adaptation and mitigation can create multiple co-benefits by simultaneously generating other ecosystem services important to human health and well-being in cities (see Figure 8.4).

8.2 Challenges for Maintaining Urban Biodiversity and Ecosystem Services

Biodiversity protection and adaptive urban ecosystem management, planning, and restoration are critical to maintain a resilient supply of climate-relevant UES in the face of global environmental change (McPhearson et al., 2014a). Globally, urban land cover is projected to increase by 1.2 million square kilometers by 2030, nearly tripling the urban area in 2000; this could result in considerable loss of habitats in key biodiversity

hotspots, including the Guinean forests of West Africa, the tropical Andes, the Western Ghats of India, and Sri Lanka (Seto et al., 2012). Mediterranean habitat types are particularly affected by urban growth because they support a large concentration of cities as well as many habitat-restricted endemic species – species that occur nowhere else in the world (Elmqvist 2013). Although urban land area globally comprises a small fraction of total land area, the impacts of urbanized land on biodiversity, ecosystem services, and other environmental impacts are wide-reaching (McPhearson et al., 2013c; Schewenius et al., 2014).

For example, expansion of urban development into the world's remaining hotspots (see Figure 8.5) for species and genetic diversity has implications for both urban and global biodiversity. These changes have downstream impacts on local ecosystem service provisioning that can feed back to influence urban climate and regional climate change. The direct and indirect effects of land-use changes outside of cities, which can include damming of rivers, water diversions, and agricultural practices, can also have effects on the capacities of ecosystems inside cities to function and produce services (Schewenius et al., 2014; Seto,

Case Study 8.2 New York’s Staten Island Bluebelt

Jack Ahern

Department of Landscape Architecture and Regional Planning, UMass Amherst

Robert Brauman

NYC Department of Environmental Protection, New York

Keywords	Urban stormwater management, urban biodiversity, blue-green network, ecosystem-based adaptation
Population (Metropolitan Region)	20,153,634 (U.S. Census Bureau, 2016)
Area (Metropolitan Region)	17,319 km ² (U.S. Census Bureau, 2010)
Income per capita	US\$56,180 (World Bank, 2017)
Climate zone	Dfa – Continental, fully humid, hot summer (Peel et al., 2007)

SUMMARY

New York faces growing impacts from climate change, an with increasing frequency of extreme weather events such as the 2012 Hurricane Sandy. The city’s Staten Island Bluebelt stormwater management practice is one of the best cases of an integrated ecosystem based adaptation (EbA) and disaster risk reduction (DRR) response wherein traditional water bodies and depressions are managed to accommodate and slow flood water. Native vegetation sites are developed by expanding, buffering, and linking with existing parks and conservation areas to form an ecological network to deliver multiple ecosystem services.

CASE DESCRIPTION

The Staten Island Bluebelt is a system of created wetlands developed since the 1990s to provide alternative, ecosystem-based stormwater management services in a rapidly developing borough of New York (NYC). The Bluebelt has become a model for providing multiple ecosystem services including stormwater management, water quality improvement, wildlife habitat provisioning, environmental education, and increased property values.

CLIMATE CHANGE ISSUES

NYC has been facing growing hazards and risks of climate change such as sea-level rise, storm surge, rising temperatures, and other related issues. Hurricane Sandy was an example of an extreme climate event. Mean annual temperature and precipitation in NYC increased 4.4°F (2.44°C) and 7.7 inches (19.55 cm), respectively, from 1900 to 2011, and sea level (at the Battery) has risen 1.1 feet (33.5 cm) since 1900. Climate models projections predict that, by 2050, the temperature will rise by 6°F (3.33°C) and precipitation by 15% (New York City Panel on Climate Change [NPCC2], 2013). NYC

is now working with academia, civil society, and others to make the city’s infrastructure and population more resilient and its infrastructure development sustainable. The high water table, poor drainage, and extensive wetlands of Staten Island challenge the development of a conventional stormwater drainage system. Cities across the world can learn from this good practice of stormwater management.

ADAPTATION STRATEGY

In 1990, the NYC Department of Environmental Protection conceptualized the Bluebelt program and began constructing stormwater best management practices (BMP) along stream and wetland corridors to attenuate routine storm flow and improve water quality and flood flow (Ryan, 2006). The Bluebelt concept had two principal goals: (1) to provide basic stormwater infrastructure and (2) to preserve the last remaining wetlands in Staten Island. Since 1995, more than 50 sites have been developed under the Bluebelt program, all of which were justified by a cost-benefit analysis comparing Bluebelt development costs with those of a conventional piped stormwater storage system. The cost-benefit analysis indicated a direct saving of US\$30 million (<http://cooper.edu/isd/news/waterwatch/statenisland>).

The Bluebelt’s principal function is to slow, store, treat, and attenuate stormwater in created wetlands and stormwater BMP in a self-regulating native ecosystem. Bluebelt facilities are designed as a “treatment train” of BMPs starting with a constructed “micropool” or fore-bay that receives stormwater from a trunk outlet. The stormwater flow then passes to an extended detention wetland where water is attenuated through contact with native wetland plants and soils. Native wetland plants sequester nutrients and add oxygen to wetland soils, facilitating the bacterial breakdown of nitrogen and phosphorous. Field stones are installed in culvert bottoms to reduce stream velocity and provide fish habitat.

Bluebelt design practices emphasize native plant species and communities including rare and near-extinct plants. Wetland plants are sourced from local nurseries or rescued from local development sites using custom excavating buckets to enable transplanting the full soil profile along with the wetland trees and shrubs. Bioengineering techniques including fascines, mats, and rolls are used to restore and stabilize slopes and stream banks with native wetland tree and shrub species. Bluebelts are constructed to intentionally include habitat “niches” with brush piles, downed trees, and boulder piles. Removed trees with roots attached are placed in the bottoms of Bluebelt ponds to create diverse microhabitats for fish and amphibians. Dead trees are left standing to provide habitat for cavity-nesting birds (Brauman et al., 2009).

Bluebelts are carefully designed to fit and complement their community context. For example, dams and bridges are built from field-stone to evoke the character of the region’s many historical bridges and dams. Bluebelt sites are selected to expand, buffer, and link up with existing parks and conservation areas, forming an ecological network to deliver multiple ecosystem services. Bluebelt trails are designed to link adjacent parks and provide direct community access for recreation. The Adopt-a-Bluebelt program has been successful in engaging community residents and environmental groups with basic maintenance tasks.

Water quality and flow monitoring by the U.S. Environmental Protection Agency found nutrient-removal rates exceeded the national standards for pollutant removal. Wildlife monitoring by the



Case Study 8.2 Figure 1 *Extended detention weir, Conference House Park, Staten Island Bluebelt.*

Photo: Jack Ahern



Case Study 8.2 Figure 2 *An aesthetic bridge connecting different ecosystems.*

Audubon Society has found a large number of breeding birds in the Bluebelt, including green herons, wood thrushes, and great-crested flycatchers. Fish passage provided by fish ladders support migratory breeds, such as the American eel, that go upstream to spawn. Mosquitoes are controlled through the Bluebelt’s constant through-flow of water that minimizes their breeding grounds as well as the support that BMP provides to populations of beneficial insects that feed on mosquitoes.

LESSONS LEARNED

The Bluebelt is a good example of a “green infrastructure” – a hybrid engineered and natural system designed to provide a suite of specific urban infrastructure and ecosystem services. It represents an example of an efficient system because of its innovations and collateral ecosystem services, including wildlife habitat provision, community recreation and education and increased property values.

Motivated by the success of this case, the Bluebelt concept is being exported to other NYC boroughs under the City’s multiple plans,

including the High Performance Infrastructure and new stormwater management plans and the NYC sustainability plans “PlaNYC 2030” and “OneNYC.” Bluebelts are also being considered to address ongoing combined sewer overflow (CSO) problems in other boroughs under the Jamaica Bay Watershed Plan. However, in other NYC boroughs, land use is more intensive and there are few existing wetlands and large areas of undeveloped land. In these boroughs, blue belts will be built on public lands, including highway verges and parks.

CONCLUSION

The Bluebelt is an effective adaptation response to effects of climate change on an urban environment. Staten Island was directly in the path of the 2012 Hurricane Sandy, and the Bluebelt demonstrated its resilience and adaptability. Although the storm surge and intense precipitation from Sandy exceeded the treatment capacity of the Bluebelt, it returned to a functional condition soon after the storm passed. The Bluebelt has saved the city more than US\$80 million in comparison with a conventional stormwater drainage system (Mayor’s Office, 2012).

Box 8.1 Urban Ecosystem Services

Urban ecosystem services (UES) refer to those ecosystem functions that are used, enjoyed, or consumed by humans in urban areas and can range from material goods (such as water, raw materials, and medicinal plants) to various non-market services (such as climate regulation, water purification, carbon sequestration, and flood control) (Gómez-Baggethun et al., 2013). The Millennium Ecosystem Assessment (MA) classified ecosystem services into four different categories: (1) provisioning services, (2) supporting services, (3) regulating services, and (4) cultural services (Convention on Biological Diversity [CBD], 2009), which have been modified and updated by The Economics of Ecosystems and Biodiversity (TEEB, 2010) project and applied to the urban context (TEEB, 2011) (see Figure 8.2). Provisioning services include the material products obtained

from ecosystems, including food, fiber, freshwater and genetic resources. Regulating services include water purification, climate regulation, flood control and mitigation, soil retention and landslide prevention, pollination, and pest and disease control. Cultural services are the nonmaterial benefits from ecosystems including recreation, aesthetic experience, spiritual enrichment, and cognitive development, as well as their role in supporting knowledge systems, social relations, and aesthetic values (Andersson et al., 2015b; Chan et al., 2011). Finally, supporting or habitat services are those that are necessary for the production of all other ecosystem services including provisioning of habitat for species, primary production, nutrient cycling, and maintenance of genetic pools and evolutionary processes (Gómez-Baggethun et al., 2013).

Table 8.1 Key abiotic, biotic, and cultural functions of green urban infrastructure. Source: Adapted from Ahern, 2007

Abiotic	Biotic	Cultural
Surface-groundwater interactions	Habitat for generalist species	Direct experience of natural ecosystems
Soil development process	Habitat for specialist species	Physical recreation
Maintenance of hydrological regime(s)	Species movement routes and corridors	Experience and interpretation of cultural history
Accommodation of disturbance regime(s)	Maintenance of disturbance and successional regimes	Provisions of sense of solitude and inspiration
Buffering of nutrient cycling	Biomass production	Opportunities for healthy social interactions
Sequestration of carbon	Provision of genetic reserves	Stimulus of artistic/abstract expression(s)
Modification and buffering of climatic extremes	Support of flora-fauna interactions	Environmental education

2013; Ignatieva et al., 2011). Moreover, the ability of species to move within and among urban landscapes is considered a key issue of biodiversity adaptation to climate change, one that suggests the need for cities to improve habitat connectivity and use green corridors for healthy, functioning urban ecosystems.

Cities and urban regions often have a perhaps surprisingly high level of biodiversity, including both native species and non-native species from around the world (Müller et al., 2010; Aronson et al., 2014). Urban species can therefore be an important component of regional and global biodiversity. Cities are often concentrated along coastlines, major rivers, and islands, which are also areas of high species richness and endemism, with many cities existing in close proximity to protected areas (see Figure 8.6) (Güneralp et al., 2013; McDonald et al., 2013). However, because expanding urban areas encompass an increasingly larger percentage of global biodiversity hotspots, it is all the more critical to safeguard urbanized biodiversity hotspots and promote ecological conservation in urban, peri-urban, and nearby rural areas.

Ecosystems are highly fragmented in urban areas, which can alter the genetic diversity and long-term survival of sensitive species. To ensure viable urban populations, urban planners and designers need to understand species' needs for habitat quality and connectivity among suitable habitat patches. For example, the connectivity of the habitat network within the urban area can play a major role for ground-dwelling animal movement, as for the European hedgehog in Zurich (Braaker et al., 2014). Understanding and planning for greater habitat connectivity through the use of green corridors is a key tool for city planners to design appropriate management and conservation strategies of urban biodiversity and to improve the resilience of species to climate change. Furthermore, it is important to understand how the impacts of climate change in cities will create risks and affect the vulnerability of urban ecosystems. The ability of ecosystems to sustain levels of biodiversity at or above the thresholds necessary for maintaining ecosystem integrity is critical to sustainable delivery of ecosystem services important for meeting urban sustainability and resilience goals (Andersson et al., 2015a).

ECOSYSTEM SERVICES: DEFINITIONS AND EXAMPLES





Ecosystem Service	Service Description	Example
PROVISIONING SERVICES: SERVICES THAT DESCRIBE THE MATERIAL OR ENERGY OUTPUTS FROM ECOSYSTEMS		
Food 	Ecosystems provide the conditions for growing food. Food comes principally from managed agro-ecosystems, but marine and freshwater systems, forests and urban horticulture also provide food for human consumption.	In Havana, Cuba (1996), a significant proportion of the urban population's food was produced within urban gardens, including 8,500 tons of agricultural produce, 7.5 million eggs and 3,650 tons of meat.
Raw Materials 	Ecosystems provide a great diversity of materials for construction and fuel including wood, biofuels and plant oils that are directly derived from wild and cultivated plant species.	Non-timber forest products such as rubber, latex, rattan and plant oils are very important in trade and subsistence—the annual global trade in such products is estimated to amount to US\$101 billion.
Fresh Water 	Ecosystems play a vital role in providing cities with drinking water, as they ensure the flow, storage and purification of water. Vegetation and forests influence the quantity of water available locally.	Estimates of the value of the services of a South African mountain fynbos ecosystem with an area of only 4 km ² indicated that water production was the biggest contributor to the total value of the system. The value was estimated to range from approximately US\$4.2M to 66.6M in 1997, according to how well the system is managed.
Medicinal Resources 	Biodiverse ecosystems provide many plants used as traditional medicines as well as providing raw materials for the pharmaceutical industry. All ecosystems are a potential source of medicinal resources.	80% of the world's people are still dependent on traditional herbal medicine (WHO 2002), while the sale of medicines derived from natural amounts to US\$57B per year.

Figure 8.3 Investing in urban ecosystems for climate adaptation and mitigation can create multiple co-benefits by simultaneously generating other ecosystem services important to human health and well-being in cities. Ecosystem services can be divided into four categories: provisioning services, regulating services, habitat or supporting services, and cultural services, with examples of each. For references in this table, see TEEB Manual for Cities (2011).

Source: Modified and adapted from TEEB Manual for Cities (2011). For references in this Figure, see TEEB Manual for Cities

REGULATING SERVICES: SERVICES THAT ECOSYSTEMS PROVIDE BY REGULATING THE QUALITY OF AIR AND SOIL OR PROVIDING FLOOD AND DISEASE CONTROL, ETC

Local climate and air quality regulation



Trees and green space lower the temperature in cities whilst forests influence rainfall and water availability both locally and regionally. Trees or other plants also play an important role in regulating air quality by removing pollutants from the atmosphere.

In Cascine Park in Florence, Italy, the urban park forest was shown to have retained its pollutant removal capability of about 72.4 kg per hectare per year (reducing by only 3.4 kg/ha to 69.0 kg/ha after 19 years, despite some losses due to cutting and extreme climate events). Harmful pollutants removed included O3, CO, SO2, NO2, and particulate pollutants as well as CO2.

Carbon sequestration and storage



Ecosystems regulate the global climate by storing greenhouse gases. As trees and plants grow, they remove carbon dioxide from the atmosphere and effectively lock it away in their tissues; thus acting as carbon stores.

Urban trees too, are important in carbon sequestration: in the United States, their annual gross carbon sequestration amounts to 22.8 million tons of carbon per year (as calculated in 2002). This is equivalent to the entire USA population's emissions in five days. This sequestration service is valued at US\$460 million per year, and US\$14,300 million in total.

Moderation of extreme events



Ecosystems and living organisms create buffers against natural disasters, thereby preventing or reducing damage from extreme weather events or natural hazards including floods, storms, tsunamis, avalanches and landslides. For example, plants stabilize slopes, while coral reefs and mangroves help protect coastlines from storm damage.

In the case of the Californian Napa City, USA, the Napa river basin was restored to its natural capacity by means of creating mudflats, marshes and wetlands around the city. This has effectively controlled flooding to such an extent that a significant amount of money, property, and human lives could be saved.

Waste-water treatment



Ecosystems such as wetlands filter effluents. Through the biological activity of microorganisms in the soil, most waste is broken down. Thereby pathogens (disease causing microbes) are eliminated, and the level of nutrients and pollution is reduced.

In Louisiana, USA, it was found that wetlands could function as alternatives to conventional wastewater treatment, at an estimated cost saving of between US\$785 to 34,700 per hectare of wetland (in 1995).

Erosion prevention and maintenance of soil fertility



Soil erosion is a key factor in the process of land degradation, desertification and hydroelectric capacity. Vegetation cover provides a vital regulating service by preventing soil erosion. Soil fertility is essential for plant growth and agriculture and well-functioning ecosystems supply soil with nutrients required to support plant growth.

A study estimated that the total required investment to slow erosion to acceptable rates in the USA would amount to US\$8.4 billion, yet the damage caused by erosion amounted to US\$44 billion per year. This translates into a US\$5.24 saving for every US\$1 invested.

Pollination



Insects and wind pollinate plants which is essential for the development of fruits, vegetables and seeds. Animal pollination is an ecosystem service mainly provided by insects but also by some birds and bats.

Some 87 out of the 115 leading global food crops depend upon animal pollination including important cash crops such as cocoa and coffee.

Biological control



Ecosystems are important for regulating pests and vector borne diseases that attack plants, animals and people. Ecosystems regulate pests and diseases through the activities of predators and parasites. Birds, bats, flies, wasps, frogs and fungi all act as natural controls.

Water hyacinth was brought under control in southern Benin using three natural enemies of that plant. Whereas the biological control project cost only US\$2.09 million in present value, its accumulated value is estimated to amount to US\$260 million in present value (assuming the benefits stay constant over the following 20 years), representing a very favourable 124:1 benefit cost ratio.

Figure 8.3 (continued)

HABITAT OR REGULATING SERVICES:

THESE SERVICES UNDERPIN ALMOST ALL OTHER SERVICES. ECOSYSTEMS PROVIDE LIVING SPACES FOR PLANTS OR ANIMALS: THEY ALSO MAINTAIN A DIVERSITY OF PLANTS AND ANIMALS.

Habitats for species



Habitats provide everything that an individual plant or animal needs to survive: food, water, and shelter. Each ecosystem provides different habitats that can be essential for a species' lifecycle. Migratory species including birds, fish, mammals and insects all depend upon different ecosystems during their movements.

In a March 2010 article, IUCN reports that habitat loss is the single biggest threat to European butterflies, and may lead to the extinction of several species. Habitat loss was said to occur most often as a result of changes in agricultural practice, climate change, forest fires, and expansion of tourism.

Maintenance of genetic diversity



Genetic diversity (the variety of genes between, and within, species populations) distinguishes different breeds or races from each other, providing the basis for locally well-adapted cultivars and a gene pool for developing commercial crops and livestock. Some habitats have an exceptionally high number of species which makes them more genetically diverse than others and are known as 'biodiversity hotspots'.

In the Philippines, an initiative to conserve local varieties of rice aided in the development of rice strains that are better adapted to local conditions - giving greater yield, a quality seed supply, and decreasing dependence on plant breeders - at a much lower cost than that of formal plant breeding.

CULTURAL SERVICES:

THESE INCLUDE THE NON-MATERIAL BENEFITS PEOPLE OBTAIN FROM CONTACT WITH ECOSYSTEMS. THEY INCLUDE AESTHETIC, SPIRITUAL AND PSYCHOLOGICAL BENEFITS.

Recreation and mental and physical health



Walking and playing sports in green space is a good form of physical exercise and helps people to relax. The role that green space plays in maintaining mental and physical health is increasingly becoming recognized, despite difficulties of measurement.

A review article examined the monetary value of ecosystem services related to urban green space, based on 10 studies, including 9 cities from China and 1 from the USA. It reported that on average, 'Recreation and Amenity' and 'Health effects' contributed a value of US\$5.882 and US\$17.548 per hectare per year respectively to the total average of US\$29.475 per hectare per year provided by the seven identified ecosystem services in the various studies.

Tourism



Ecosystems and biodiversity play an important role for many kinds of tourism which in turn provides considerable economic benefits and is a vital source of income for many countries.

In 2008 global earnings from tourism summed up to US\$944 billion. Cultural and eco-tourism can also educate people about the importance of biological diversity.

Based on the amounts of money people spent on travel and local expenditure in order to visit Coral reefs in Hawaii, it was estimated that the value associated with these reefs amounted to US\$97 million per year. This implies that reef tourism resulted in significant income generation for individuals, companies, and countries.

Aesthetic appreciation and inspiration for culture, art and design



Language, knowledge and the natural environment have been intimately related throughout human history. Biodiversity, ecosystems and natural landscapes have been the source of inspiration for much of our art, culture and increasingly for science.

Prehistoric rock art of southern Africa, Australia, and Europe, and other examples like them throughout the world, present evidence of how nature has inspired art and culture since very early in human history. Contemporary culture, art and design are similarly inspired by nature.

Spiritual experience and sense of place



In many parts of the world natural features such as specific forests, caves or mountains are considered sacred or have a religious meaning. Nature is a common element of all major religions and traditional knowledge, and associated customs are important for creating a sense of belonging.

In the example of the Maronite church of Lebanon, the church committed to protecting a hill in their possession, comprising rare remainders of intact Mediterranean forest, independent of scientific and legal arguments, because this was in line with Maronite culture, theology and religion.

Figure 8.3 (continued)



Figure 8.4 Investing in urban ecosystems and green infrastructure can provide multiple co-benefits. This shows a cultural co-benefit of urban and peri-urban trees through tapping sugar maple (*Acer saccharum*) trees in Pound Ridge, New York. Maple sugar tapping represents a seasonally occurring peri-urban and urban food production ecosystem service that has long-standing cultural traditions in many northern countries.

Photo: Timon McPhearson. Adapted from Andersson et al., 2015b

8.2.1 Current Effects of Climate Change on Urban Biodiversity and Ecosystems

All urban ecosystems will experience the effects of climate change. Additionally, many cities are located in geographic areas that are especially vulnerable to both existing and projected climate hazards, such as coastal flooding, landslides, and extreme events. Climate change is impacting a broad spectrum of urban ecosystem functions, biodiversity, and ecosystem services (Rosenzweig et al., 2011; Solecki and Marcotullio, 2013; UN-Habitat, 2011). Urban ecosystems are already under general stress from development, pollution, and direct human use (Elmqvist et al., 2013), and climate change variability poses additional challenges for urban species and ecosystems. For example, in a comprehensive review of the potential impacts of climate change on urban biodiversity in London, Wilby and Perry (2006) highlight the importance of four threats to biodiversity in the city: competition from non-native species, pressure on saltmarsh habitats from rising sea levels, drought effects on wetlands, and changing phenology of multiple species as earlier springs occur more frequently (Hunt and Watkiss, 2011).

The UHI effect in cities can change the reproductive and population dynamics of animals. Insect life cycles and migration

Case Study 8.3 The Serra do Mar Project, Baixada Santista Metropolitan Region (BSMR), São Paulo State

Oswaldo Lucon

São Paulo State Environment Secretariat

Keywords	Resettlement, biodiversity protection, climate resilience, urban ecology, floods, landslides, ecosystems
Population (Metropolitan Region)	1,664,136 (IBGE, 2015)
Area (Metropolitan Region)	2,405 km ² (IBGE, 2015)
Income per capita	US\$14,810 (World Bank, 2017)
Climate zone	Af – Tropical rainforest (Peel et al., 2007)

A partnership of the Inter-American Development Bank (IDB) and the São Paulo State Government, the Serra do Mar and Mosaics System Recovery Program has been recognized as an international standard for resettling communities in disaster-prone, ecologically sensitive areas. *Mosaics* are sets of protected areas located nearby or juxtaposed to each other. Their main purpose is to promote integrated and participatory management of their components, respecting different categories of management and conservation objectives. Mitigation strategies comprise halting deforestation, reforestation, and wastewater treatment. Adaptation strategies are based on the resettlement of populations living in landslide- and flood-prone areas. The Program started in the city of Cubatão and part of the



Case Study 8.3 Figure 1 Brazilian Atlantic Forest, before (left) and currently (right).

Baixada Santista Metropolitan Region (BSMR) of 2,405 square kilometers (IBGE, 2015). Topography varies from cliffs (700 m) to plains (average 3 m above sea level). The Atlantic Forest is a UNESCO Biosphere Reserve, one of the planet’s biologically richest regions and also one of the most endangered. Overexploitation and biome devastation have resulted in only 7% of the Brazilian Atlantic Forest being preserved in fragments of more than 100 acres (see Case Study 8.3 Figure1).

In the Southeastern State of São Paulo, the Atlantic Forest is concentrated in the *Serra do Mar*, squeezed between the coastal BSMR (nine cities, 1.6 million people) and the São Paulo Metropolitan Region (19 cities and 20 million people). The 133,000 square kilometers of the Serra do Mar State Park cover twenty-four municipalities in the state. Additionally, three mosaics (Paranapiacaba, Jureia-Itatins, and Jacupiranga) allow for buffer zones between urban and native preserved areas. Despite having been reduced and highly fragmented,

the Atlantic Forest is habitat to more than 20,000 plant species – a wealth of diversity greater than that found in North America (17,000 species) and Europe (12,500 species). Out of the native plant species, 8,000 are endemic; that is, native species that only exist in Brazil (IAD and São Paulo State Government, 2009). Degradation of the forest had its origins in the construction of roads. In the highly industrialized city of Cubatão, settlements on hillsides (*bairros-cota*) invaded areas belonging to the Serra do Mar State Park. Illegal occupations harmed not only the Park, but also created several hazards to its inhabitants: landslides, floods, road accidents, and freshwater contamination (see Case Study 8.3 Figure 2).

In the first stage, the São Paulo Government contributed 65% of the US\$470 million budget, and the IDB allocated the remaining 35%. Geotechnical studies mapped and classified risk areas.

Potential damage to dwellings and their residents was estimated, considering their positions and distances to critical slopes plus the degree of building vulnerability (construction pattern and level of urban consolidation). A joint analysis of these criteria established a mapping of risk sectors, with hotspots defined (IDB and São Paulo State Government, 2013; see Case Study 8.3 Figure 3). In 2007, new settlements were halted (“frozen”) through supervision of the Military Police, with protective measures for the Park that included preventing deforestation, fires, and the capture of wild animals and extraction of plant species, and monitoring of the various sectors of the *bairros-cota* to prevent their expansion (IDB and São Paulo State Government, 2013).

A total of 7,388 irregular households were identified, with around 7,760 families and 7,843 buildings. The resettlement program was



Case Study 8.3 Figure 2 Settlements on hillsides (*bairros-cota*) invaded areas belonging to the Serra do Mar State Park, creating several hazards to its inhabitants, including landslides, floods, road accidents, and freshwater contamination.



Case Study 8.3 Figure 3 Designated risk areas.

followed by an environmental education program, an enrollment process, sealing buildings, commissioning basic housing and urbanization projects, obtaining environmental licenses, conducting public hearings, and negotiating with the IDB for co-financing. Benefits include improved living conditions for around 3.2 million people in the surrounding area, an increase to 60,000 visitors per year in the Park, improved biodiversity, improved water quality, strengthened management and protection of conservation units (an additional 20,000 hectares of Atlantic Forest; recovery of 1,240 hectares of State Park), and lowered disaster risk, plus more sustainable sources of income.

More than 5,000 families living in at-risk or protected areas have been resettled and assisted with housing and upgraded infrastructure. Living in new structured communities, they have also benefited from professional training programs for construction professionals, gardeners, and nurserymen to work on the reforestation of the reclaimed areas. The second phase of the program aims at assisting approximately 25,000 families with resettlement or infrastructural upgrading. Building improvements included two or three types of houses with diversified typologies, accessibility for the disabled, preservation of significant green areas, and improved urban infrastructure. Family assistance combines social, cultural, economic, and environmental aspects. Resettlement has brought innovations

that enabled families to feel sufficiently engaged before and after moving from their homes, including the choice of one of fifteen housing options. Housing units were not donated, and leaving a house where one had lived for a long time is not an easy decision, even if it means moving to better conditions. Therefore, for families who live in rural or peri-urban areas, other methods have been developed.

To anchor all actions, synergy among institutions has proved decisive. In 2009, the state joined the United Nations Environment Program (UNEP)'s Sustainable Social Housing (Sushi) initiative for building sustainable social housing for low-income populations. A pilot neighborhood (Residencial Rubens Lara) in Cubatão City has been developed and today is recognized by the UNEP as a replicable model for other countries. In 2012, the Serra do Mar Social and Environmental Recovery Program earned the Greenvana GreenBest award, the highest distinction conferred in Brazil for environmental initiatives. The Serra do Mar Program went beyond the limits of the City of Cubatão. It now covers the whole of the Atlantic Forest of São Paulo, extending throughout the Park (north and south of the state) to the Jureia-Itatins territory and the Units for Marine Conservation. The extended program is called Serra do Mar and the Atlantic Forest Mosaics System Social and Environmental Recovery Program (CDHU, 2012; São Paulo, 2013).

patterns have been well-documented, with changes in the life cycle of certain insects having already occurred in response to urban warming (Parmesan and Yohe 2003; Parmesan, 2006). Butterfly species in Ohio, for example, appear to have shifted when they fly in response to urban warming. Some native butterfly species appear to be at risk due to the shortening of their flight periods (Kingsolver et al., 2013). In Raleigh, North Carolina, the abundance of the gloomy scale butterfly (*Melanaspis tenebricosa*) increases with increases in impervious surfaces that create warmer forest temperatures and therefore drive increased reproduction rates, thus contributing to greater population growth for this urban forest pest (Dale and Franck 2014). This suggests that urban trees could face greater herbivory in the future as a consequence of the increased fitness of some herbivorous arthropods under warming scenarios. However, more research is needed to generalize these results to other urban areas.

Although other ecological and socioeconomic factors are affecting vegetation in urban areas, many of the non-native invasive species colonizing cities originate in warmer areas and are benefiting from changing climate conditions (Sukopp and Wurzel, 2003). In mountain regions, climate is already causing changes in vegetation structure and diversity (Theurillat and Guisan 2001; ICIMOD, 2009). The response of trees to extreme climatic events may be species-specific. For example, in Dresden, a study of oak trees showed that *Quercus petraea* and *Q. rubra* are better adapted to warm and dry conditions than are *Acer platanoides* and *A. pseudoplatanus* (Gillner et al., 2014).

Trees have been perhaps better studied than other taxonomic groups in urban areas. Urban trees experience multiple forms of stress including heat stress, low air humidity, and soil drought. Rapid climate change can have a significant impact on the distribution and biology of trees. In Philadelphia, climate change

is influencing the biology of urban tree pathogens and pests. Results from a recent study indicate that the future climate in Philadelphia will become less optimal for multiple tree species since major pests and diseases are likely to become more problematic (Yang, 2009).

Comparing urban and rural species has yielded a useful understanding of urban biodiversity responses to changing climate. Woodall et al. (2010) compared tree species compositions in northern urban areas to tree compositions in forestland areas. They found that some tree species native to eastern U.S. forests of southern latitudes have been planted or are present in northern urban forests, indicating the tolerance of southern species in northern urban ecosystems. Although urbanization and climate change can both profoundly alter biological systems, scientists often analyze their effects separately. Recent studies are beginning to look at these impacts on organisms simultaneously to better understand how multiple simultaneous stressors might affect species, but more research is needed (Kingsolver et al., 2013).

8.2.2 Projecting Impacts of Climate Change on Urban Biodiversity and Ecosystems

Projecting impacts of climate change on the distribution of species is complex, with many factors to consider including dispersal ability, species interactions, and evolutionary changes (Pearson and Dawson 2003; Gilman et al., 2010; Urban et al., 2012). Still, future climate change in cities, when combined with additional urban stressors such as short-lived climate pollutants, land use change, and direct human impacts is expected to pose difficult challenges for urban species and ecosystems. Maintaining adequate levels of biodiversity and managing urban ecosystems to ensure a resilient supply of critical ecosystem services that are necessary for expanding urban populations may




Figure 8.5 As urban areas expand into some of the most sensitive biodiversity areas globally, urban biodiversity conservation and ecological planning will have an increasing impact on global biodiversity and ecosystems and the services they provide to both urban and rural residents. Urban areas with large populations in 1970, 2000, and 2030 (projected) are shown in green as examples of urban expansion in global biodiversity hotspots (in blue). The cities shown have a projected population of more than 8 million in 2030, according to UN World Urbanization Prospects 2014.

become increasingly challenging in the future as climate change intensifies its effects on cities. Which ecosystems will be most affected in the near and longer term may be signaled by current species' responses to climate change (Parmesan, 2006; Gillner et al., 2014). The risks and vulnerabilities associated with climate change in urban ecosystems are likely to vary with temporal and spatial scale and nature of change (e.g., chronic vs. acute), although in general they are expected to increase over the next several decades (Solecki and Marcotullio, 2013).

We present here new regionally downscaled climate projections using the Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report (AR5) scenarios for forty global cities spanning small, large, and megacities in multiple contexts including coastal and inland cities in the Global North

and Global South in the 2020s, 2050s, and 2080s (see Figure 8.7a, 8b) (see Chapter 2, Urban Climate Science, and ARC3.2 Annex 2, Climate Projections for ARC3.2 Cities). Projections for both temperature and precipitation show wide variation in cities around the world, with temperature generally increasing and precipitation both increasing and decreasing depending on location. Effects on ecosystems will vary considerably from city to city, and therefore it is not possible to suggest general management or planning approaches. Instead, decision-makers in cities and urban areas will need to take into account locally relevant climate projections combined with data on sensitive species or ecosystems to develop plans and adaptive management strategies to safeguard urban ecosystems and the benefits they provide for climate adaptation and mitigation (as well critical co-benefits for human well-being). These downscaled climate projections suggest that urban planning, policy, and management must pay close attention to decisions and actions involving urban ecosystems that may be directly impacted by uncertain climate futures. Climate change in cities is already having significant impacts on urban biodiversity and ecosystems. Further impacts of rising temperatures and increasing or decreasing precipitation suggest increasing ecological impacts over time, with concurrent affects on the ability of urban ecosystems to provide nature-based solutions for building climate resilience in cities.

8.3 Climate Change Hazards, Risks, and Vulnerabilities for Urban Ecosystems

When combined with socioeconomic changes, there are multidimensional vulnerabilities affecting biodiversity and urban ecosystems. Heat stress, inland and coastal flooding, droughts, cyclones, fire, and extreme rainfall pose risks to urban ecosystems, populations, and economies (Revi et al., 2014). Massive land conversion from natural ecosystems to a built environment exposes urban landscapes to loss of biodiversity, flash floods, droughts, and pollution while urban sprawl and poor urban design further threaten urban biodiversity (Munaung et al., 2013; Revi et al., 2014). Recent studies demonstrate how climate change is reinforcing urban ecosystem vulnerability through unsustainable development, agricultural land conversion, and degradation of ecosystem services that affect the ability of ecosystems to meet urban climate adaptation and mitigation goals (Satterthwaite et al., 2007; UN-Habitat, 2009, 2011).

8.3.1 Climate Hazards and Risks

Urban climate hazards are defined as the climate-induced stressors or drivers that affect urban ecosystems. Examples include elevated temperature, changes in precipitation patterns, sea level rise, and the build-up of short-lived climate pollutants such as black carbon (see Figure 8.7), as well as changes in the frequency and intensity of extreme events such as storm surge, flash floods, heat and cold waves, and wild fires (UNEP, 2011). The cascading effects of climate change can have both direct and indirect effects on biodiversity and ecosystems. Climate change




Figure 8.6 Urban areas are expanding into protected areas in all parts of the world. Figure shows (a) urban extent and (b) percentage of total urban extent within a distance of, from top to bottom, 10, 25, and 50 kilometers of protected areas (PAs, e.g., national parks) by geographic region circa 2000.

Source: Adapted from McDonald et al., 2013; Güneralp and Seto, 2013

Case Study 8.4 Ecosystem-Based Climate Change Adaptation in the City of Cape Town

Pippin M. L. Anderson

Department of Environmental and Geographical Science, University of Cape Town, Cape Town

SUMMARY

Cape Town is adapting to growing urban climate change vulnerability and impacts. The city, with its rich biodiversity and unique ecosystems, historically used hard engineering measures to reduce growing flood and storm surge risks. However, in recent years, the role of ecosystem services is being recognized and included in urban climate change adaptation plans. Recent initiatives by the city administration to identify and spatially map urban ecosystem services (UES), in particular in relation to the bionet (network of green open spaces) map, to establish critical connectivity corridors suggest a good start in mainstreaming climate change in urban development planning and environment conservation.

CASE DESCRIPTION

Cape Town, with an area of 2,460 square kilometers and a population of approximately 3.7 million has close to 38% low-income households, indicating high poverty incidence. The city’s population also has a high disease burden due to the high prevalence of HIV and tuberculosis. More than 58% of the adult population has below a

Keywords	Ecosystem-based adaptation, disaster risk reduction. flood water, wetlands, stormwater management
Population (Metropolitan Region)	3,740,025 (City of Cape Town, 2015)
Area (Metropolitan Region)	2,461 km ² (City of Cape Town, 2015)
Income per capita	US\$12,860 (World Bank, 2017)
Climate zone	Csb – Temperate, dry summer, warm summer (Peel et al., 2007)



Case Study 8.4 Figure 1 Iconic Table Mountain of Cape Town viewed from the Durbanville Conservation Area.

Photo: Pippin Anderson

high school education, and 16.9% of the population is unemployed. The city is characterized by urban sprawl and rapidly expanding informal poor settlements on the lowland areas that are known as the Cape Flats. The increasing demand for housing continues to place a burden on city authorities and on remnant urban biodiversity.

Cape Town is located in the Cape Floristic Region, the smallest and most diverse floral kingdom on earth: the region hosts almost 9,000 plant species on 90,000 square kilometers – some 44% of the flora of the subcontinent on a mere 4% of its land area. Of approximately 3,350 indigenous plant species within the metropolitan boundary, 190 are endemic to the city that also hosts 19 of 440 National Vegetation Types (Cilliers and Siebert, 2012). The process of urbanization has significantly contributed to the erosion of local biodiversity, putting further stress on eleven nationally recognized critically endangered vegetation types in the city. The City is host to 83 mammal, 364 bird, 60 reptile, 27 amphibian, and 8 freshwater fish species. The lowlands historically hosted the greatest vegetation-type and floral diversity, and the majority of this has been lost to urban settlements. Some 450 of these indigenous plant species are listed as threatened or near-threatened, and 13 species are known to be extinct.

Remnant natural ecosystems are highly fragmented, with little connectivity. Fire is used as a management tool in a burning rotation of 10–15 years, which poses a management challenge in an urban setting, threatening both property and life. Introduced invasive plant species suppress indigenous biodiversity and yield high fuel loads that, under a rising temperature regime, lead to hotter and more dangerous fires.

CLIMATE CHANGE VULNERABILITY AND IMPACTS

Climate change is occurring faster in South Africa than the global average. Mean annual temperatures have increased faster than the global average during the past 50 years. Extreme rainfall and drought events have also increased in frequency (Ziervogel et al., 2014). Urban areas are particularly vulnerable due to stormwater

surge, flooding, uncontrolled fire, and coastal erosion. The Cape Town region is likely to face significant climate change risks with predicted increases in temperature in all seasons, reductions in rainfall, greater evaporation, more intense and frequent winds, and greater coastal erosion and storm surge with changes in the frequency and intensity of extreme weather events. Increased rainfall intensity will exacerbate flooding, especially in high water table areas on the Cape Flats. Flooding is exacerbated due to the canalized nature of rivers where natural vegetation buffers have been removed.

Cape Town is a water-scarce area. Current climate change predictions suggest increased rainfall variability with associated future increases in periods of drought and water shortages. Climate change predictions suggest hotter, more frequent, and runaway fires. Cape Town, with its 307 kilometers of coastline, is at threat from climatic hazards such as sea level rise and increased storm surge.

ADAPTATION STRATEGY

The City has adopted an integrated water resource management (IWRM) approach that includes demand-side water management. Acknowledging the role of invasive plant species in reducing water availability, the government public works program seeks to train and employ unskilled and unemployed labor to clear invasive vegetation, producing positive outcomes in biodiversity, social benefits, and water yield.

Adaptation measures to increased flood risk include both engineering and ecological solutions that includes the creation of retention ponds and resilient infrastructure, regular drain cleaning, better disaster warning systems, the decanalizing of rivers, and the restoration of riparian vegetation to vulnerable areas. However, engineering solutions get less attention mostly due to high costs and flood disaster-relief funding structures.

Fire, an ecologically necessary measure to promote indigenous flora, is being used more judiciously. The intensity and season of firing



Case Study 8.4 Figure 2 *Controlled burning of urban vegetation.*

are being regulated to have positive implications for biological processes of recruitment and regeneration. Fire regimes during periods of drought, higher wind speeds, and generally greater climate variability are being used strictly for assured biodiversity and employment generation. Government public works – “Working for Water” and “Working on Fire” programs – are used to reduce large fuel loads and minimize runaway fires. These programs train firefighters in ecological fire management using higher public safety protocols. In general, these programs, set up to address various environmental and social issues, have proved to be an important vehicle for generating adaptive capacity and change in the face of threats posed by climate change.

DRIVERS

The City Administration has taken a number of measures to adapt to climatic changes and mitigate threats. Historical measures, such as sea embankments to protect infrastructure, are now recognized as extremely expensive to maintain and as sometimes ineffective. Acknowledging the high costs of these engineering measures, the City is employing more ecosystem-based approaches including the protection and restoration of extensive wetlands sites that can

absorb large volumes of water and dissipate wave energy (ICLEI, 2012). Efforts are on to restore dune vegetation and to open paths to improve sand supply to these mobile systems that have frequently become cut off due to hard engineering solutions employed in the past. These ecosystem-based adaptation (EbA) measures are providing green employment and thus contributing to the City’s poverty reduction goal.

IMPACT AND LESSONS LEARNED

These multipronged adaptation approaches have drawn involvement from multiple stakeholders and worked to create better impacts and synergy. The City is trying to secure the establishment of a “bionet” – a network of green open spaces – that would serve to improve biodiversity areas by allowing for greater flexibility and opportunity for species conservation, provide vegetated areas for water infiltration, and reduce flooding and storm surge impacts. The role of ecosystem services will become critical in the face of climate change. The initiatives by City government to identify and maintain ecosystem services and a biodiversity corridor suggest a good start in mainstreaming climate change in urban development planning and environment conservation.

also has significant economic and human impacts that can extend from infrastructure and built environment sectors to natural ecosystems (Frumkin et al., 2008; Keim, 2008; Hallegatte et al., 2010; Ranger et al., 2010; Solecki and Marcotullio, 2013). For example, in cities with diminishing precipitation, the vegetated cover of green roofs may face drought risks. Increased exposure due to rising populations and growth of human settlements in flood- and landslide-prone areas exacerbate climatic hazards as well as socioeconomic risks, thus emphasizing the sensitive interactions among climate urban ecosystems, and communities.

8.3.1.1 Thermal Hazards

Changing temperature regimes (see Figure 8.7a) can have both direct and indirect effects on the organisms that live in

cities and the ecosystem services that they provide. At the individual and species population level, many physiological processes such as photosynthesis, respiration, growth, and flowering of plants are affected by changing temperature. Elevated temperature can affect growth and reproductive rates either positively or negatively for plants (Hatfield, 2011), while also inducing a range of landscape-level impacts on biogeochemical cycles and watershed hydrology (Suddick et al., 2012). Warming conditions in New York, for example, have led to changes in tissue chemistry in tree seedlings relative to cooler, non-urban settings, resulting in more rapid shoot growth but reduced root mass (Searle et al., 2012). Higher temperatures can also lead to increased physiological stress on wildlife, affecting their behavior and reproduction (Marzluf, 2001) (see also Section 8.2.1).

Table 8.2 Effects of urban climate and environment on urban agriculture. Source: Adapted from Wortman and Lovell, 2013

Drivers of plant production	Compared to rural areas	Observed positive effects	Observed negative effects	Resulting impact on urban crop yield	Expected future dynamics of drivers
Length of growing season	7 to 8 days longer	Potential of double cropping systems		Higher	Increase
Time to flowering	Earlier		Risk of asynchrony between timing of flowers and pollinator presence	Lower	Increase
CO₂ concentration	Higher	Increased photosynthesis rate in many vegetable crops (C ₃ plants)		Higher	Increase
Temperature	Higher	Increased photosynthesis rate up to threshold	Decreased photosynthesis rate (in case of extreme temperature), increased irrigation water demand	Higher, lower	Increase
Wind speed	Lower	Reduced plant mechanical damages	Increased leaf gas exchanges	Higher	?
Vapor pressure deficit	Higher (less air humidity)		Greater plant transpiration, moisture stress, reduced photosynthesis rate, reduced rainwater infiltration in soil, lower soil moisture	Lower	?
Ground-level ozone concentration	Higher (sometimes lower)		Decreased photosynthesis rate, lower root-to-shoot ratio, premature leaf senescence	Lower-higher	Increase
NO₂ concentration	Higher	More efficient nitrogen nutrition	Delayed flowering, accelerated plant senescence	Higher-lower	Increase
Soil water infiltration	Lower		Higher moisture stress	Lower	Increase-decrease

8.3.1.2 Drought Hazards

More intensive increases or decreases in precipitation can lead to significant water-related urban hazards including drought and severe water shortages (IPCC, 2013). Reductions in precipitation can be exacerbated by warming temperatures, which increase water losses to evapotranspiration driven by hydrological alterations from surface-water diversion and groundwater extraction (Pataki et al., 2011). Increased frequency and duration of droughts exacerbated by warming can also increase evapotranspiration (Leipprand and Gerten, 2006). Increased evapotranspiration reduces water availability and groundwater resources, often leading to increased salinization and water stress affecting both the quality and quantity of water for plants, with negative consequences on floral and faunal biodiversity and productivity (Alberti and Marzluf, 2004). For example, projected drought conditions in Manchester, England, are likely to reduce the cooling services provided by grasslands, which may increase the local UHI and wild fires (Gill et al., 2013). Current drought in California is affecting drinking water supplies and is also having dramatic effects on peri-urban agriculture; this has led to historic water conservation measures to deal with drought stress. Drought affects both street trees and urban parklands and will likely have cascading effects on herbivores, soil fauna, and other components of urban biodiversity, as well as effects on urban

residents through decreased water availability affecting livability (Wilby and Perry 2006; Gillner et al., 2014).

8.3.1.3 Flood Hazards

More frequent and increased precipitation (see Figure 8.7b) can lead to significant urban flood hazards. Flash floods, in addition to damaging critical infrastructure and directly impacting the lives of urban dwellers, also are harmful to urban water supplies and drainage systems and can have lasting negative impacts on ecosystems (IPCC, 2013). Increasing extreme precipitation events in combination with land-cover changes and increased frequency of tropical cyclones and subsequent altered water flow in urban watersheds is likely to result in an increased incidence of flooding in many cities (Depietri et al., 2012; IPCC, 2013). Flood hazards include the short-term impacts of the force of moving water (e.g., flash floods), inundation, and drowning, which cause longer-term impacts resulting from sediment movement (erosion and deposition), soil processes, and the distribution of pathogens that precipitate negative public health impacts (ICIMOD, 2012; Teegavaerapu and UNESCO, 2012; Wisner et al., 2003; Walker et al., 2008). For cities along rivers and coastlines, rising sea levels and increasing storm surges will increase urban flooding as well (Mosely 2014). Coastal flooding due to sea level rise can




Figure 8.7 a) Here we show projected average temperature and precipitation for forty global cities in 2020, 2050, and 2080 (see Chapter 2, Urban Climate Science, and ARC3.2 Annex 2, Climate Projections for ARC3.2 Cities) and major habitat types. Cities represent a range of small, large, and megacities in the Global North and Global South, including inland and coastal cities. Temperature (8.7a) (in °C) and precipitation (8.7b) (% change) projections are based on thirty-three Global Climate Models and two Representative Concentration Pathways (RCP4.5 and RCP8.5) downscaled from regional to city spatial extent. Changes are relative to the 1971–2000 base period. The time slices are the 30-year periods on which the projections are centered (e.g., the 2020s is the period from 2010 to 2039).

lead to increased salinization and reduced groundwater recharge (Chan et al., 2011; IPCC, 2013), which can decrease habitat quality for biodiversity.

Climate change in cities will lead to increased precipitation in some places and decreased precipitation in others (see Figure 8.7a). In cities projected to receive increased precipitation,

increased discharge into surface waters will have ecosystem consequences. For example, urban development affected the ability of watersheds in Baltimore, Maryland, to retain nitrogen, and urban watersheds showed increased sensitivity to climate variation (Kaushal et al., 2008). Loss of this urban ecosystem function in Baltimore (nitrogen retention) led to increased nitrogen downstream, with negative consequences for the ecology

Case Study 8.5 Jerusalem Gazelle Valley Park Conservation Program

Naomi Tsur

Former Deputy Mayor, Jerusalem Green Fund

Helene Roumani

Jerusalem LAB Coordinator, Local Action for Biodiversity

Keywords	Biodiversity, urban planning, development, Jerusalem mountain gazelle, ecosystems
Population	839,000 (CIA World Factbook, 2015)
Area	125 km ² (www.worldcat.org, 2017)
Income per capita	US\$36,190 (World Bank, 2017)
Climate zone	Csa – Temperate, dry summer, hot summer (Peel et al., 2007)

CASE DESCRIPTION

The historic city of Jerusalem is also a well-known place for its rich natural heritage of Biblical flora and fauna that has developed as an integral part of the city landscape. The city is a significant habitat for half a billion birds since it lies on one of the most important global bird migration routes following the course of the Great Rift Valley. The credit for creating this rich ecosystem and wealth of biodiversity within a city lying in a water-scarce area goes to the Gazelle Valley Park Conservation Program (Krasny, 2015).

The major issue of climate change faced by the city is shortage of water and the threat of desertification. Temperatures in the Middle East region are not rising as fast as in other parts of the world, but the region is already experiencing weather extremes and the process of desertification is on the rise. Although rainfall has increased, so has evaporation. The impact of climate change on the region’s natural flora and fauna is still mild mainly because of their historical adaptive capacity to withstand moisture stress and high temperatures. However, future predictions are that, due to extreme heat and water stress, plants and animals will have difficulty surviving.

CLIMATE CHANGE ADAPTATION STRATEGY

The City of Jerusalem has assumed the responsibility for improving and maintaining its unique desert and hilly ecosystems to preserve its floral and faunal biodiversity in the face of increasing climate change stresses. In 2009, Jerusalem joined the ICLEI–Local Governments for Sustainability/Local Action for Biodiversity (ICLEI/LAB) Network to further pursue sustainable development measures.

In the context of Jerusalem’s LAB Legacy project for the International Decade of Biodiversity, Jerusalem has established the Gazelle Valley Conservation Program to protect and restore one of the city’s unique biodiversity areas and to plan the development of a park for both wildlife preservation and local recreation at the site. The area has recently been designated as an urban nature park – a local model of sustainable development.

The Gazelle Valley is situated on a sixty-acre undeveloped tract of land in southwest Jerusalem, between two residential neighborhoods and closed in by major roadways. After being used for agricultural purposes during the 1960s and 1970s, the land, a rich wildlife habitat, was left as open space while the surrounding urban area continued to develop. The mountain gazelle (*Gazella gazella*), an indigenous species particularly prevalent in this part of the Jerusalem hills, has been roaming the valley and sustaining itself on its natural resources since ancient times. It is also the site of ancient terraces with orchards that still bear fruit.

In the late 1990s, a residential plan was established for the Gazelle Valley, threatening to destroy the gazelle habitat and remove a vital open space in the city. The Jerusalem branch of the Society for the Protection of Nature (SPNI) opposed the development plan, citing that it was a reversal of established urban planning principles. Local residents and activists joined SPNI to launch a campaign to save the Valley.

The Gazelle Valley Citizen Action Committee was thus formed. Understanding the need for a comprehensive plan, the Committee, together with SPNI, commissioned an alternative plan focusing on conservation and restoration of the site’s unique biodiversity. After 10 years of rigorous grassroots opposition, the city decided to withdraw the residential plan and designate the Gazelle Valley a natural heritage



Case Study 8.5 Figure 1 Mountain gazelles, examples of revived urban biodiversity urban biodiversity.

site. In addition, the conservation plan was approved by the Local Planning Committee in 2009, marking the first time that local authorities approved a development plan initiated by residents. This civil society initiative for environmental protection in Jerusalem was also a significant victory for the environmental movement in the region.

CLIMATE CHANGE ADAPTATION

The development of the Gazelle Valley Park in Jerusalem plays an important role in the city's promotion of climate change adaptation. Water conservation is a significant aspect of the park design. Apart from the need to regulate the drainage basin, water features prominently in the plan as a vital natural resource for sustaining the local biodiversity. In addition, regulation of existing water systems is being planned to enhance the beauty of the site and serve to attract visitors. The plan includes a series of runoff collection pools that will have the capacity to store 20,000 cubic meters of rainwater and seepage. In addition, a runoff filtration system is also planned for sedimentation of solids in water entering the park. To control seasonal flood zones, the valley's natural irrigation system will be rehabilitated, facilitating the restoration of the site's ancient agricultural terraces. In order to

prevent erosion and control channel flow, two gravel-lined streams will be dug in alignment with the local topography. Proper rainwater management will not only create a buffer zone between the conservation area and the adjacent recreational area, it will also help mitigate climate change effects and the effects of increased urbanization around the Valley.

IMPACT AND LESSON LEARNED

The park is expected to serve local residents and visitors with a public activity core (differentiated from the animal habitat), including pedestrian and bike paths, gazelle observation points, a bird watching route, agricultural gardens, and an educational visitor center. The Gazelle Valley Conservation Program in Jerusalem demonstrates that through proper planning practices, conservation efforts in an urban setting can facilitate both climate change adaptation and promote efficient ecosystem management. In the case of Jerusalem, it is anticipated that this effort will also produce an effective interface between biodiversity and human activity. The city government is taking the lead in mobilizing stakeholders in steering this green adaptation project.



Case Study 8.5 Figure 2 *Gazelle Valley Park, Jerusalem City.*

and economy of the Chesapeake Bay. On the other hand, freshwater wetlands with reduced hydrologic inputs could become even further water-limited, with negative effects on both ecosystem services and biological diversity (World Bank, 2015). The modification of climate within and around cities combined with increasing drought stress from decreased precipitation illustrates how climate change will affect many urban ecosystems worldwide (UN-Habitat, 2011).

8.3.1.4 Hazards Related to Shifting Species Distributions

Species movement in response to climatic regime shifts has already been well-documented (Parmesan and Yohe, 2003; Parmesan 2006; ICIMOD, 2009; Porter et al., 2013). Organisms

at the edge of their distributions may decline as temperature or other climate conditions shift outside their physiological tolerances. For example, plant and animal species may shift northward seeking cooler temperatures following climate shifts, meaning that cities in the tropical and subtropical belts may lose species faster (Gonzalez et al., 2010; Grimm et al., 2013). Changing climate may also affect the introduction of new species in urban ecosystems by reducing noninvasive and native species while favoring weedy and urban-adapted species (Kendle and Forbes 1997; Booth et al., 2003; Heutte et al. 2003). Warming cities (see Figure 8.7a) may find new problems with invasive species and pests that had formerly been limited by cold conditions. The most adaptive species in an era of changing urban climate are likely to include more weeds, pests, and invasive species, such as

the introduced Burmese python in Florida (IPCC, 2014). Species that are highly specialized and heat sensitive may be threatened with local extinction driven also by an inability to move to new areas as urban development expands.

The distribution of pathogens is also likely to shift with changing climate, with consequences for both resident organisms and the ecosystem functions they provide. For example, climate change is likely to influence the distribution of the mosquito *Aedes aegypti*, the primary urban vector of dengue and yellow fever viruses (Eisen et al., 2014) (see Chapter 10, Urban Health). Ultimately, changes in species distributions are expected to modify the ecological interaction networks in cities and have the potential to promote invasive species, which could accelerate the loss of urban biodiversity (Nobis et al., 2009; Kendal et al., 2012).

It is important to recognize that none of these hazards and risks operates in isolation. For example, changes in CO₂ concentrations may or may not amplify the impacts of changes in precipitation, temperature, or other climate hazards on urban vegetation (Zavaleta et al., 2003), suggesting the need for further research to better understand critical feedbacks in the urban system. Thus, integrating all of the ecosystem processes and recognizing that there are critical feedbacks among ecological, built, and social components of urban systems will yield a more thorough understanding of climate risks to urban biodiversity and ecosystems.

8.3.2 Urban Ecosystem Vulnerability

Vulnerability may be considered a lack of resilience or a reduction in adaptive capacity (Tyler and Moench, 2012). However, the complexity of urban ecosystems is characterized by vulnerability along multiple dimensions. Urban ecosystems share many of the same types of climatic vulnerabilities as non-urban ecosystems. However, urban ecosystems are also exposed to a number of unique stressors and therefore they experience greater exposure to hazards such as a high concentration of pollution, the inherent role of non-climate stressors, and the UHI phenomenon (Farrell et al., 2015). The extent of human conversion of the landscape and anthropogenic inputs of materials, energy, and organisms are all greater in cities, which can affect climate vulnerability in

a variety of ways (Fitzpatrick et al., 2008; Loarie et al., 2008). Rapid urban growth and local landscape dynamics⁵ contribute to national, regional, and global-scale climate change driven by elevated rate of greenhouse gas (GHG) emissions, radiative forcing⁶ of non-greenhouse gases, and alteration of rainfall patterns by short-lived climate pollutants (e.g., black carbon, tropospheric ozone, and methane) (Cervený and Balling, 1998; Pielke et al., 2002; Parmesan, 2006; UNEP, 2011).

In the urban ecosystem context, exposure to multiple stressors is a real concern, particularly in developing countries where socioeconomic and political drivers along with climate variability play important roles (Leichenko and O'Brien, 2008). However, there are very few studies that have assessed the multidimensional nature of urban ecosystem vulnerability important for planning appropriate adaptation measures (IPCC 2007, Williams et al., 2008). Assessing the vulnerability of urban ecosystems to climate change is critical to include as part of urban planning, policy, and design processes that intend to ensure sustainable delivery of ecosystem services into the future (McPhearson et al., 2015a) (see Chapter 5, Urban Planning and Design).

Vulnerability of urban ecosystems can be assessed at multiple levels within the urban system, including for the individual organism (e.g., physiological health and reproductive success of humans, plants, and other biota), populations, and for larger landscapes (e.g., land use and land cover, biogeochemical cycling) (UNEP-WCMC, 2009; Vignola et al., 2009; Kalusmeyer et al., 2011; Violin et al., 2011). Most studies to date examine vulnerability at the species level or, in some cases, landscape level. Williams et al. (2008) and Glick et al. (2011) have developed species-level vulnerability assessments in which they define “species vulnerability as a function of climate change–related impacts and the adaptive capacity of the species.” However, given the strong connections among urban, peri-urban, and rural landscapes, it is important to assess combined and connected cumulative effects of exposure and sensitivity to climate change. Kalusmeyer et al. (2011) argue that assessing vulnerability at landscape level is cost effective and a more useful tool for decision-makers than, for instance, than, for example, vulnerability assessments focused on single species (see Box 8.2).

Box 8.2 Ecosystem Vulnerability

Klausmeyer et al. (2011) used a vulnerability assessment tool for climate change impacts on biodiversity using landscape-scale indicators in California. This method allows biodiversity managers to focus analysis on the species likely to be most vulnerable and to decide on the best adaptive strategies to reduce vulnerability to climate change. Based on results, the authors recommended that state biodiversity

managers focus on minimizing current threats to biodiversity (9% area), reducing constraints to adaptation (28%), reducing exposure to climatic changes (24%), and implementing all three (9%). In 12% of the high-vulnerability areas, current conservation goals have to change; in remaining areas, no additional actions are required. This tool can also help to identify adaptation measures focused on endangered species only.

⁵ Landscape dynamics is a concept of landscape equilibrium highlighting the spatial and temporal scaling of disturbance regimes and their influence on equilibrium/non-equilibrium dynamics in a particular landscape (Pielke et al., 2002).

⁶ “Radiative forcing is a measure of the influence a factor has in altering the balance of incoming and outgoing energy in the Earth-atmosphere system, and is an index of the importance of the factor as a potential climate change mechanism” (IPCC, 2001).

8.4 Adaptive Capacity and Urban Ecosystem Resilience

The adaptive capacity of species in urban landscapes is a function of ecology, physiology, and genetic diversity (Kalusmeyer et al., 2011; Williams et al., 2008). The adaptive capacity of an urban ecosystem is also the degree to which system dynamics can be modified to reduce risk. Traditionally, adaptive capacity focused on human actors and institutions, but, in the context of urban biodiversity and ecosystems, nonhuman actors, behavior, species interactions, and human–ecological interventions are also important. For example, human-induced adaptive capacity could include planting species that are more tolerant of higher temperatures and droughts. Nonhuman-derived adaptive capacity could include natural processes that change ecosystem components rapidly for organisms like insects populations persisting despite changing climate. Adaptation measures such as introducing green infrastructure (e.g., urban green spaces, constructed wetlands, agricultural land in outlying flood-prone areas) can reduce thermal loads and flood hazards and improve water and air quality for vulnerable biota (Depietri et al., 2012) (see also Case Study 8.2, Staten Island Bluebelt, and Case Study 8.5, Jerusalem Gazelle Park). In addition, cities are dependent on urban and peri-urban ecosystems for food production, water provision, and air-quality regulation, meaning that the adaptive capacity of a specific urban area depends at least partially on local-to-regional considerations (Tyler and Moench, 2012).

Resilience to climate change is a growing priority among urban decision-makers. Improving resilience will require transformations in social, ecological, and built infrastructure components of urban systems (Tyler and Moench, 2012; Ernstson et al., 2010) (see Chapter 1, Pathways to Urban Transformation). Urban ecosystems are important components when building urban resilience through their ability to absorb climate-induced shocks and ameliorate the worst effects of extreme climate events (McPhearson et al., 2015a). However, disturbances of sufficient magnitude or duration, such as prolonged drought, can push biodiversity and ecological relationships beyond safe thresholds for reliable production of ecosystem services and may require new approaches to land-use planning and adaptation that focus on building ecological resilience (Folke et al., 2004).

Cities are increasingly seeking to enhance adaptive capacity of urban ecosystems through, for example, green infrastructure, including urban agriculture, landscape conservation, green roofs, green walls, and other green and open spaces that conserve ecosystem values and functions (Kremer et al., 2016a). Building urban parks and other green spaces and adding vegetation strips to densely built neighborhoods can help reduce thermal hazards, manage stormwater, and enhance health benefits, thus enhancing climate change resilience (e.g., in Rio de Janeiro). From a climate and resource-efficiency perspective, the spatial configuration of green spaces is particularly important to mitigate the UHI effect and to conserve water and energy use. Cities with a combination of a high percentage of green areas, high edge

density (distribution of the green space), and high patch density (number of patches per unit area) can more effectively respond to climate extremes such as heat waves and heavy precipitation (European Environment Agency [EEA], 2015; Maimaitiyiming et al., 2014). This suggests that policy and urban planning should ideally prioritize connected green corridors of critical mass rather than a multitude of fragmented green spaces; nevertheless, the total percentage of green space independently is likely most impactful for climate resiliency and, in practice, is often more feasible to create.

Many cities are vulnerable to the hazards associated with climate change as a function of their location (UN-Habitat, 2011). For example, cities are disproportionately distributed along coasts and major rivers, which increases their vulnerability to floods and storm surges. Urban ecosystem managers planning species- and landscape-level adaptation often have multiple goals such as protecting land, restoring habitat, encouraging compatible lands uses, and reducing fragmentation (Heller and Zavaleta, 2009). Building resilient urban ecosystems therefore needs flexible, modulating, and safe-to-fail approaches that can adapt to uncertainty and extreme climate events such as typhoons and hurricanes (e.g., Hurricane Sandy in New York, 2012) (Tyler and Moench, 2012). Also, greater coordination and networks among governance structures that manage local ecosystems and urban biodiversity, including cemeteries, golf courses, urban parks, and neighborhood gardens, would strengthen ecosystem functioning as well as the associated and essential social-ecological engagement (Ernstson et al., 2010).

8.4.1 Interactions between Social and Ecological Infrastructure

The vulnerability of urban ecosystems and biodiversity is intrinsically linked to human activities that drive urban system dynamics. The urban population, with its resource consumption and waste-generation activities, the built infrastructure system (buildings, transportation infrastructure, utilities), and the direct and indirect modifications to the landscape (e.g., changes in vegetation, water courses and storage, microclimate) all create a distinct set of vulnerabilities for the systems and biota embedded in cities (Alberti, 2015). These vulnerabilities are manifested at multiple spatial scales. At the very local level, the altered microclimate and hydrology of a city street will affect the ecosystem services generated by local trees, wildlife, and microbes. Within larger ecosystems embedded in cities, such as remnant forests and urban agriculture and wetlands, the direct effects of human activities and infrastructure need to consider both local and landscape-level management to reduce hazards exposure and risks simultaneously at multiple levels.

8.4.2 Adaptive Management of Vulnerable Urban Biota

The multidimensional nature of urban vulnerability impacts urban biodiversity components including the diversity of

plants, animals, and microbes within city boundaries. These groups are all influenced by environmental changes associated with both urbanization and human management. City managers should support both biological communities that have persisted since before urban development (e.g., remnant forest patches, indigenous wildlife that have adapted to urban conditions) and novel communities that depend on human inputs (e.g., pests, deliberately or accidentally introduced species) (Aronson et al., 2014). For example, cities create novel ecosystems and habitats outside their natural biome, such as warm subway tunnels in cold regions, lakes and ponds in arid areas, and dry soils

in humid areas that contribute to increased biodiversity levels often observed in urban areas compared to surrounding ecosystems (McKinney, 2002).

Urban biodiversity vulnerability can be mediated by direct and indirect human management of habitats. The response of indigenous species in remnant ecosystems is affected by regional climate shifts, local ecological dynamics, and the local impact of the city itself (e.g., augmented warming, altered water resources, direct human impact). These urban influences can be moderated by direct human management that reduces their exposure and

Case Study 8.6 Medellín City: Transforming for Life

Leonor Echeverri

Administrative Department of Planning, Medellín City Council

Keywords	Urban development; transportation; adaptive urban planning; resilient infrastructure, ecosystems
Population (Metropolitan Region)	3,731,000 (Alcaldía de Medellín, 2015)
Area (Metropolitan Region)	1,152 km ² (Alcaldía de Medellín, 2015)
Income per capita	US\$13,910 (World Bank, 2017)
Climate zone	Am – Monsoon-influenced humid subtropical (Peel et al., 2007)

Medellín is an inspiring example of sustainable and innovative urban development with good governance, community participation, and business partnerships. City leaders can take some credit for transforming the city into a vibrant, socially cohesive, and more environmentally resilient city through initiating adaptive and flexible urban planning strategies with effective implementation. The positive impact of mass transportation, green spaces, and equitable benefit sharing resulted from citizen participation, stakeholder involvement, and government support for urban development. The effective use of social networks and good communication by city leaders sustained community support.

CASE DESCRIPTION

Medellín, located in mountainous Aburrá Valley, is a Colombian city with a history of sustainable urban development processes. Many of the poor communities living on the mountainous slopes were challenged for safety and access to essential city services. City leadership has since provided public safety, security, easy mobility, access, amenities, and opportunities. Medellín also developed affordable mass transport systems – the world’s first cable car system – the Metrocable – and also a Metro to address both access and pollution problems. Today, Medellín is famous for its social cohesion, business-friendly environment, people and environment-centric city governments and a high quality of life. How Medellín transformed itself from a city with high socioeconomic challenges to one described as “a great inspiration to other cities facing similar issues” can be attributed to Medellín’s bold, visionary leadership, which

encompassed diverse stakeholders to deliver a series of small-scale but high-impact, innovative green urban projects (Eveland, 2012).

SOCIOECONOMIC AND ENVIRONMENTAL ISSUES

By the 1970s, Medellín demographically had grown by almost twenty-fold – from around 60,000 in 1905 to more than 1 million – to become the second-largest city of Colombia. A large number of poor were living in precarious socioeconomic and ecological conditions, suffered exclusion, and were struggling with a high cost of living. Medellín entered a cycle of decline in its economic base, which led to a consolidation of a segregated, unequal, and conflict-ridden society and degrading ecosystem. By the early 1980s, Medellín faced a host of social and economic upheavals that led to government failure. In response, the city unleashed social mobilization processes that constituted the genesis for a collective construction of a new vision of urban development, one that led to political and strategic processes that began major change and development pathways in the city. The community responded with significant efforts toward collective dialogue in which a broad cultural and pedagogical process laid the groundwork for civic and citizen-led projects. This included environmentally sensitive urban development planning and program implementation.

ADAPTIVE CHANGE PROCESS

Affordable mobility played an important role in achieving equitable connectivity between urban and rural sections in Medellín (Moreno et al., 2013) As well as setting a process of forming neighborhoods in response to functional interests and a population demanding specific interventions, expanding the city’s services to include green spaces throughout the city improved its greening index, making the city more climate resilient (Green City Index [GCI], 2010).

Milestones were conceived during the 1980s and 1990s as the Strategic Plan of Medellín and the Metropolitan Area 2015. This generated a broad and pluralistic project of continuity and consistency in a society that was in crisis. A participatory process was developed for sustainable development that became a foundation for environmentally friendly policies and practices.

In 1995, the Metro became operational – a point of origin for the Integrated Mass Transit System that linked physical, institutional, virtual, sustainable, and environmental modes of mass transit with efficiency and effectiveness. The Metro system serves the current as well as future transportation needs of all inhabitants. This has helped Medellín to minimize its ecological footprint and protect biodiversity and ecosystem services in spaces freed-up by the Metro.

A joint exercise between government and development agencies has been proposed to assess the vulnerability dynamics of the territory, along with implementing sustainable alternatives to mitigate climate impacts in both urban and rural areas of the Valle de Aburrá. With a holistic adaptation approach, criteria are enforced to ensure the security of both the people and the ecosystem within the city's territory. The Integrated Transport System of the Aburrá Valley (SITVA), the Inventory of Greenhouse Gases, the Environmental Classrooms Program of Integrated Solid Waste Management, Linear Parks and Ecological Corridors, Best Practices of Sustainable Consumption and Production, the More Forests to Medellín project, and Integral Water Management among others have positively contributed to the improvement of indicators of an adaptive urban system. Linear parks in particular help the city protect itself from storms and increased pollution.

IMPACTS AND LESSONS LEARNED

The city's Green Belt encourages conditions and opportunities for integral human resources development in the transition zone between urban and rural regions. The Green Belt is important as a way to regulate city expansion into sensitive ecosystems and has helped to conserve and protect natural habitat.

The Medellín River Park is another urban renewal project, connecting the city with efficient mobility, public space, and environmental interventions. Engineering and urbanism work hand-in-hand so that the city's rivers can form the structural axis of civic life.

This ongoing process of transformation has shown that it is possible to build a community-driven, environmentally friendly project in a city. The city's development plan is based on a territorial focus on its urban-rural areas and contains a systemic view to overcome inequities. This has inspired bottom-up planning processes and public-private partnerships to find innovative alternatives.

Medellín's Home for Life initiative recognizes that a participative society and good governance are combined in an institution that seeks equity as a result of political and social rationality. Here, the urban development goes beyond different forms of land use and integrates a combined human-environment urban ecosystem framework. The lessons learned in these efforts will prove useful in confronting the daunting challenges of adapting to climate change.

CONCLUSION

The main driver of Medellín's transformation has been city government's efforts to be inclusive, fair, participatory, and environmentally sound in urban development governance. These approaches transformed Medellín into a model of sustainable urban livability and earned it the 2014 Lee Kuan Yew World City Prize Special Mention. Medellín aspires to continue advancing as an innovative and intelligent city, and hopes to facilitate the exchange of experiences and the advancement of collective knowledge among cities and their inhabitants. To promote sound green design and appropriate policies embracing multidimensional development, building resilient rules, regulations, capacity, and citizen's participation have been the key factors.



Case Study 8.6 Figure 1 Parque de Los Pies Descalzos (Barefoot Park)

Photo: Municipality of Medellín and Development Plan 2012–2015: Medellín, a home for life

sensitivity, for example by eradicating pest organisms or creating conservation programs for rare or endangered and endemic species. The Gazelle Valley Park Conservation Program is an example of how climate change adaptation can be combined with biodiversity conservation through ecosystem management (see Case Study 8.5).

Another example is how integrated urban water management can reduce the vulnerability of urban ecosystems and biodiversity (see Case Studies 9.5 and 14.B, Rotterdam). Management of water resources for drinking and sanitation, as well as the hazards associated with water (flooding, landslides, etc.), can alter water flow and storage for the benefit of urban plants. Management of urban hydrological systems through improved greening can decrease the vulnerability of urban ecosystems. For example, during drought periods, a small share of water resources may be reserved as environmental flow for use by plants and animals, thus allowing ecological systems such as forests, wetlands, and streams to survive and maintain adaptive capacity. While drought may affect an entire region, urban ecosystems where water resources are well managed can reduce the impact of such climate-driven water stress, but only provided that urban ecosystem management activities are part of a larger system-level urban resilience plan.

8.5 Ecosystem-Based Adaptation and Nature-Based Solutions

In the urban context, healthy ecosystems can replace or complement often expensive “hard” or engineered infrastructure (e.g., sea-walls, dykes or embankments for river control, and shelters). EbA⁷ and similar nature-based solutions have been

widely recognized as “soft,” safe-to-fail, and often less expensive approaches to climate resilience that values and uses ecological services for adaptation (Huq et al., 2013; CBD, 2009). EbA approaches can generate numerous co-benefits and indirect ecological and social benefits to both non-urban and urban communities in ways that support urban transitions to sustainable, livable communities (UNFCCC Secretariat, 2011, 2013; UNEP, 2012; Huq et al., 2013; Zandersen et al., 2014). The contribution of green infrastructure to EbA in the form of urban parks, avenue plantation, and urban forestry can also provide small levels of GHG mitigation by storing carbon in soils and vegetation. Multiple co-benefits are also expected from the integration of climate change adaptation and biodiversity conservation measures (ICLEI-Africa, 2013; UNEP, 2009). For example, although not a direct goal of climate adaptation, city green spaces have been shown to have important societal co-benefits including but not limited to lower crime rates, reduced level of stress, enhanced cognitive capacities, and improved public health (Troy et al., 2012; Demuzere et al., 2014). The Singapore Case Study (Case Study 8.7) illustrates how EbA can be integrated with disaster risk reduction strategies.

Urban ecosystem services may play an increasingly vital role as cities grow in population size and contain larger senior age cohorts than any other settled topography. The 2003 heat wave in Europe resulting in more than 70,000 deaths can be seen as an early warning of more severe climatic conditions to come, with climate change being viewed as a new public health threat (Petkova et al., 2014). In the United States, heat is the greatest weather-related cause of human death because increasing temperatures above 90°F (32°C) aggravate air pollution and ozone levels and result in greater health risks, including respiratory illness (e.g., asthma) and heart attacks, particularly

Box 8.3 Urban and Peri-Urban Agriculture and (Agro) Forestry for Climate Change Adaptation and Mitigation

Marielle Dubbeling

RUAF Foundation, Leusden

The IPCC’s AR5 (IPCC, 2013) projects that, due to climate change, there is likely to be a loss of food production and productive arable lands in many regions. Cities with a heavy reliance on food imports and the urban poor will be significantly affected. Adaptation options and local responses mentioned in the report include support for urban and peri-urban agriculture, green roofs, local markets, enhanced social (food) safety nets, and the development of alternative food sources, including inland aquaculture (University of Cambridge and ICLEI, 2014).

Urban and peri-urban agriculture have long been recognized as a potential food security and income strategy (Zezza and

Tasciotti, 2010; De Zeeuw et al., 2011; FAO, 2014; Porter et al., 2014). However, its potential contribution as a climate change adaptation and, to a lesser extent, mitigation strategy has only been more recently studied and acknowledged (Lwasa, 2013; Dubbeling, 2014; Lwasa and Dubbeling, 2015). Because a clear framework and tools for monitoring the contributions of urban and peri-urban agriculture and forestry (UPAF) to climate change mitigation and adaptation was not available until recently, the potential to integrate UPAF into city climate change plans was limited.

A recent (2012–2014) collaboration between UN-Habitat and the Resource Centers on Urban Agriculture and Food Security (RUAF) Foundation-International aimed to respond to these gaps by (1) enhancing the awareness of local authorities and

⁷ Ecosystem-based adaptation is an approach to planning and implementing climate change adaptation by considering ecosystem services and their uses for human well-being (MEA, 2005).



Box 8.3 Figure 1 Multifunctional design of urban greenways in Bobo Dioulasso (Burkina Faso).

Source: F. Skarp

other stakeholders involved in urban climate change programs, land use, agriculture, and green spaces regarding the potentials (and limitations) of UPAF for climate change adaptation and mitigation; and (2) assisting interested cities and other local actors to integrate urban agriculture into local climate change and land-use policies and strategies and to initiate pilot actions that “showcase” replicable urban agriculture models. At the same time, RUAF Foundation and the Climate Development Knowledge Network (CDKN) developed and tested a monitoring framework in an attempt to quantify the impacts of UPAF on climate change adaptation and mitigation in participating cities.

One of the partner cities, Bobo-Dioulasso, in Burkina Faso, is characterized by increasing urbanization resulting from industrial and economic growth, as well as from the return of migrants following an internal crisis in Ivory Coast. The city is built up in housing blocks with square open urban lots between them. The municipality is trying to preserve these lots (called greenways) for greening (agroforestry) and multifunctional (recreation and urban agriculture production) activities as part of their parks and gardens program and climate change adaptation strategy.

Satellite images and remote sensing data were used to quantify the effect of land uses on land-surface temperatures (LSTs). A comparison of 1991–2013 data showed that LST differences between urban and peri-urban areas increased approximately 6% a year. The study also showed that mean LSTs over a 10-year period were consistently cooler (0.3°C) in the three specific green infrastructure areas analyzed than in adjacent urbanized areas (Di Leo et al., 2015). This may have important effects on human well-being.

In addition, the greenways will contribute to increased infiltration and retention of stormwater, with a reduced runoff coefficient of 4%, thus possibly reducing flood risks in periods of intensive rainfall. Monitoring has also shown that from a first harvest cycle (August–October 2013) from open fields can contribute to at least 6% of the monthly food expenditures of the households involved in the project (RUAF Foundation, 2014).

In the same way, these urban agriculture greenways contribute to a more permanent availability of home-produced food for these households. Such increased diversification of food and income sources helps to increase the resilience of poor households, which are generally vulnerable to increases in food prices (Dubbeling, 2014). Furthermore, preservation of green infrastructure is highly relevant because municipalities in Africa, as elsewhere, regularly encourage infill developments and higher housing densities that lead to the reduction or loss of green spaces and gardens.

In the period 2013–2014, the municipality of Bobo-Dioulasso decided to (1) install and institutionalize a municipal committee for the future management of the greenways; (2) draft and adopt a technical statute for the greenways promoting their productive and multifunctional use; and (3) adopt a set of specifications applicable to the exploitation of the greenways. The draft legal texts were submitted to and adopted by the Environment and Local Development Commission of the Municipality in January 2014. On March 26, 2014, the proposal to install the municipal committee was unanimously adopted by the municipal council. A provision of €20,000 was made in the 2013–2014 municipal budget to cover the functioning and activities of the Greenway Management



Box 8.3 Figure 2 *Rehabilitated paddy areas with vegetables growing on raised bunds in Kesbewa, Sri Lanka.*

Photo: Janathakshan

Committee and to support maintenance of the existing productive greenways as well as their replication on other lots (RUAF Foundation, 2014).

In comparison to Bobo-Dioulasso, where increasing urban temperatures and urban heat islands are the main predicted climate change impacts, the city of Kesbewa, Sri Lanka, has to deal not only with increasing temperatures but also with more intense rainfall and regular flooding (see Box 8.3 Figure 2). Kesbewa is a medium-sized, rapidly expanding city located 25 kilometers from the capital Colombo. Kesbewa city used to be characterized by a large presence of agricultural and rice-producing fields, the latter in lower-lying areas and flood zones. Much of the agricultural activity has been abandoned due to rice production from the north of the country being more economically viable and due to sale of land for urbanization. The rapid filling and conversion of the paddy lands to residential and commercial areas has significantly altered natural water flows and drainage. Coupled with an increase in average rainfall and heavy rainfall events, this has resulted in recurrent flooding and related damage to infrastructure, utility supply, and the urban economy in some parts of Kesbewa (CDKN, 2014).

It was for this reason that the Ministry of Agriculture, Western Province, with support of UN-Habitat, RUAF, and the local non-governmental organization Janathakshan, decided to implement a pilot project on rehabilitating abandoned paddy lands by promoting the production of traditional varieties of salt-resistant paddy rice that fetch good market prices combined with the growing of vegetables on raised beds. By re-establishing the flood regulation and ecosystem services of these areas, this strategy will not only contribute to reducing flood risk but

also to increasing urban food production and income generation for farming households. Support for urban agriculture as a flood risk or stormwater management strategy was also taken up in cities like Bangkok after the 2012 flooding (Boossabong, 2014) and in New York (Cohen and Wijsman, 2014).

Support for this program and its expansion is being institutionalized in policy uptake at different levels. At the local level, the preservation of agricultural areas and flood zones is included in the Kesbewa Urban Development Plan. At the provincial level, urban agriculture is now considered a climate change adaptation strategy for the province. The current Climate Change Adaptation Plan 2015–2018 of the Western Province of Sri Lanka (Ministry of Agriculture, 2014) now specifically includes action lines regarding the expansion of urban and peri-urban agriculture and agroforestry, the management of paddy lands as a flood risk reduction strategy, and the reduction of food miles by promoting localized production. And, at the national level, prescribed land use for low-lying urban and peri-urban rice fields now allows for the new production model as part of the revised “Paddy Act” (RUAF Foundation, 2014).

More localized food production may also have positive impacts on reducing energy use and emissions related to food transport (cold) storage, and packaging. In both Kesbewa and Rosario, Argentina, urban consumption patterns and food flows were analyzed and scenarios developed to calculate the potential impacts of increased local food production. Assuming similar production systems would be applied in both distant rural (current production locations) and peri-urban areas for production of the main consumed



Box 8.3 Figure 3 Horticulture production in Rosario's greenbelt.

vegetables in Rosario (potato, squash, beans, and lettuce), CO₂ emissions related to such food consumption would be reduced by about 95%. Analysis of production capacity in Rosario's peri-urban areas shows that such local production is feasible.

In response, the Municipality of Rosario has already zoned – in its urban development plan – an additional 4,000 hectares of its remaining greenbelt for vegetable production and has, in collaboration with the Province of Sante Fe, started a pilot program to support horticulture producers using ecological production techniques and opportunities for direct marketing. A first agreement was signed with the Association of Hotels and Restaurants for this purpose (Dubbeling, 2015).

The amount of food that can actually be produced in urban and peri-urban areas was more recently the subject of study in Almere, the Netherlands, and in Toronto, Canada. A 2012

scenario study done in Almere found that 20% of total food demand (in terms of potatoes, vegetables, fruits, milk, and eggs) projected for a future population of 350,000 inhabitants can be produced locally, within a radius of 20 kilometers of the city. More than 50% of the needed area is devoted to animal production (grass and fodder). By replacing 20% of the food basket with local production in Almere while at the same time promoting fossil fuel reduction in production, processing, and cooling through the use of renewable energy sources, energy savings (363 TJ) would add up to the equivalent of the energy use of 11,000 Dutch households. Savings in GHG emissions (27.1 Kt CO₂ equivalent) would equal carbon sequestration in about 1,360 hectares of forest or the emissions of 2,000 Dutch households. The largest savings are due to reduction in transport, replacing fossil fuel use with renewable energy sources (solar, wind energy); use of excess heat from greenhouses; and replacing conventional with organic – or ecological – production (Jansma et al., 2012).

among urban dwellers. High GHG emissions compound temperature levels, leading to forecasts of even higher U.S. summer temperatures and health concerns in coming years (Kenward et al., 2014). As climate change becomes increasingly viewed in the context of public health, cities that incorporate green infrastructure in urban planning and the built environment, and that safeguard local biodiversity, will optimize their urban ecosystems services for temperature mitigation, thus strengthening climate resiliency as well as improving quality of life (Santiago Fink, 2016).

8.6 Urban Agriculture and Forestry

With increasing urbanization, climate change, and growing urban demand for food, cities need to address the triple challenge of climate change mitigation and adaptation, as well as the provision of basic services, including food, to vulnerable residents. Barthel et al. (2013, 2015) suggest that urban agriculture production in its many forms has been supporting urban resilience throughout the history of urban development. The examples here

show that urban and peri-urban agriculture and forestry may provide helpful strategies to address this triple challenge (see Box 8.3). The future upscaling of these interventions will need new urban design concepts and the development of local and provincial climate change action and city development plans that recognize urban agriculture as an accepted, permitted, and encouraged land use. The involvement of the subnational (e.g., provincial) government is key to addressing agriculture and land-use planning at larger scales (outside municipal boundaries), facilitating access to financing, and developing the regional policies that must accompany city-level strategies (Dubbeling, 2014).

8.7 Ecosystem-Based Mitigation Strategies

Urban areas are likely to face the most adverse impacts of climate change due to high concentration of people, resources, and infrastructure (Revi et al., 2014). Climate change mitigation is therefore required to reduce the sources and enhance the sinks of GHGs, especially carbon. Combining green infrastructure and EbA may increase urban CO₂ sinks (Rogner et al., 2007), although estimates for different kinds of green infrastructure remain contested (Pataki et al., 2011). Urban land-use changes have significant impact on GHG emissions and carbon sequestration, as well as on albedo, which plays an important role in radiative forcing. New and updated urban plans warrant the inclusion of both climate change resilience measures as well as long-term mitigation strategies that need to be supported by metrics and decision-support tools that demonstrate GHG reductions; land use and transportation as well as green infrastructure indicators are needed (Condon et al., 2009).

Integrated urban planning that incorporates a multidisciplinary perspective to target schemes that also support increased use of green infrastructure, forest restoration, and other EbA approaches can help advance sustainable urban development while reinforcing climate mitigation and enhancing the quality and quantity of UES (RUAF, 2014; Ecologic Institute, 2011; Georgescu et al., 2014). For example, incorporating green infrastructure in urban design, especially in warmer climates, can potentially reduce the use of air conditioning, cause significant energy savings, and therefore indirectly reduce GHG emissions (Alexandri and Jones, 2008; Georgescu et al., 2014).

8.8 Cross-Cutting Themes

8.8.1 Urban Planning and Design

Designing, planning, and managing complex urban systems for climate resilience and human health and well-being require ecosystems to be resilient to effects of climate change and be able to sustainably and reliably provide critical ecosystem services over time (McPhearson et al., 2014b). Urban planning and design are key processes that determine the quantity, quality, and accessibility of urban residents to UES (see Chapter 5, Urban Planning and

Design). Urban development often replaces natural elements with built and impervious surfaces, which can degrade and eliminate ecosystems, natural processes and flows (e.g., water and nutrients cycles), and biodiversity (Alberti, 2008; Colding, 2011; Novotny et al., 2010). Impervious surfaces also exacerbate climate-related problems such as the UHI effect, flooding, and other stormwater management concerns. To counter these trends, ecosystem-based approaches in urban planning and design practices are emerging. Ecosystem-based approaches can include urban green infrastructure in ways that enhance ecosystem services and restore native biodiversity. In a growing number of cities, local communities and city planners are collaborating to create new green spaces and improve existing ones using GIS and other holistic spatial planning tools and technologies (Pickett and Cadenasso 2008).

Over the past few decades, “ecocities” and “green cities” theories began to emphasize the importance of ecosystems within cities and in linked rural areas as a way to provide important ecosystem services to city residents (Yang, 2013). Innovative urban planning theories such as Ecological Design (Rottle and Yocom, 2011), New Urbanism, Sustainable Urbanism (Farr, 2008), Ecological Urbanism (Mostafavi and Doherty, 2010), Agricultural Urbanism (De La Salle and Holland, 2010), Landscape Urbanism (Waldheim, 2007), Green Urbanism (Beatley, 2000), Biophilic Urbanism (Beatley, 2009), Ecocities (Register, 2006), and Ecopolises (Ignatieva et al., 2011) emphasize ecological restoration and connected multifunctional green infrastructure in dense, compact cities. These new approaches in urban planning are beginning to prioritize walkable and mixed land uses, emphasizing designs that cater to the needs of people and other living things (Register, 2006). In this way, urbanizing areas can start to facilitate climate mitigation and adaptation as co-benefits with efforts to reduce waste and consumption (Register, 2006).

Sustainable urban design seeks to maximize the quality of the built environment and minimize impacts on the natural environment, transforming impervious areas into high-performance landscapes (McLennan, 2004). Inter- and transdisciplinary, collaborative and strategic urban planning and design, based on restoration and reconnection of green areas at different scales, can offer numerous benefits (Breuste et al., 2008; Colding, 2011;

Box 8.4 Key Messages: Cities Biodiversity Outlook

1. Rich biodiversity can exist in cities.
2. Biodiversity and ecosystem services are critical natural capital.
3. Maintaining functioning urban ecosystems can significantly enhance human health and well-being.
4. Urban ecosystem services and biodiversity can help contribute to climate change mitigation and adaptation.
5. Ecosystem services must be integrated into urban policy and planning.

Source: Adapted from Secretariat of the Convention on Biological Diversity, 2012

Novotny et al., 2010; McDonald and Marcotullio, 2011; Pauleit et al., 2011; Ignatieva et al., 2011; Ahren, 2013). For example, urban planning and design that promotes habitat connectivity through linkages or clustering of landscapes, parks, and green infrastructure can increase the provision of multiple ecosystem services such as recreation, stormwater management, and biodiversity preservation (Colding, 2011). More recent approaches to urban green infrastructure design also acknowledge ecosystem disservices (see Box 8.5), the need to account for disservices as well as tradeoffs and synergies in biodiversity, and different ecosystem services (Von Döhren and Haase, 2015; Gomez-Baggethun et al., 2013; Kronenberg, 2015).

In both urban and non-urban contexts, climate change is associated with the increased frequency and intensity of extreme events and accelerated loss of urban biodiversity (Thomas et al., 2004). Adapting to urban climate change in the face of an uncertain magnitude of risk, vulnerability, and impacts means that urban planners should have both short- and long-term adaptation options for which a constant flow of information and knowledge is critical. Ongoing assessment of the state of urban ecosystems and ecosystem services across multiple scales and functions can support the planning and design of interconnected urban social-ecological systems (Kremer et al., 2016a).

8.8.2 Equity, Environmental Justice, and Urban Ecosystem Services

Human and nonhuman vulnerabilities are intimately intertwined at the urban scale, and the most vulnerable (including both human and nonhuman) species lack the power and capacity to respond to climate change impacts (Steele et al., 2015) (see Chapter 6, Equity and Environmental Justice). From an environmental justice⁸ perspective, the quantity, quality, and accessibility of urban ecosystems and their services is unevenly distributed across urban populations, with the poor and minorities often disproportionately affected by environmental hazards and ecosystem disservices and lack of access to essential ecosystem services (Pham et al., 2012; McPhearson et al., 2013a).

For example, the location, structure, and quality of urban parks present a long-term environmental justice challenge. Access to parks provides ecosystem services benefits such as recreation, physical activity, public health, aesthetic value, education, and sense of place. Historically, it has been demonstrated that the urban poor were often forced to leave their homes to create space for the creation of urban parks (Taylor, 2011). More recent research shows that the health and well-being of minorities and low-income populations are affected by the lack of access to high-quality, large, urban parks (Boone et al., 2009; Loukaitou-sideris and Stieglitz, 2002; Miyake et al., 2010) and other kinds of green spaces – such as urban vacant lots – that produce social and ecological benefits (McPhearson et al., 2013). A recent study in Bogotá, Colombia (Escobedo et al., 2015) identified marked

Box 8.5 Ecosystem Disservices

Although urban planning and design is increasingly embracing urban ecosystems as cost-effective design solutions, it has yet to deal with the emerging knowledge of how ecosystems can also create negative impacts on human well-being, known as “ecosystem disservices.” Although green infrastructure provides a wide range of services, they also generate disservices (Gomez-Baggethun et al., 2013; McPhearson, 2014; Döhren and Haase, 2015) that are important to take into account in urban planning and management.

For example, green roofs have value in improving the quality of runoff by reducing pollutant release (Dunnnett and Kingsbury, 2008), but some studies have noted the negative effect of the roofing materials used on the quality of runoff water due to chemicals or metal compounds (Bianchini and Hewage, 2012) and also on air quality by the emission of volatile organic compounds and nitrogen oxides (Kaye, 2004).

Similarly, urban trees can produce pollen that negatively affects allergy sufferers and may affect asthma rates in cities. Working closely with local ecological experts when developing green infrastructure will be important to understand the inherent tradeoffs to maximize ecosystem services and minimize disservices associated with particular species or species assemblages (McPhearson et al., 2014; Döhren and Haase, 2015).

inequalities in ecosystem services provision by urban trees. The poorest socioeconomic stratum had the lowest tree and crown size, whereas the wealthiest stratum had the largest tree attributes.

Minorities and the poor are also more likely to use urban biodiversity directly as a source of livelihood and thus are more impacted by the effect of climate change and pollution on natural resources such as fisheries and urban agriculture, especially in low- and middle-income countries (Corburn, 2005; National Environmental Justice Advisory Council, 2002). It will be important to consider the spatial distribution of environmental justice in planning and decision-making on policies related to ecosystem services. For example, the location of new green infrastructure can improve environmental justice by locating natural spaces and elements in proximity to otherwise underserved populations. The opposite may be true if new green infrastructure is located at the expense of such populations, where, for example, gentrification processes together with new green space development increase the cost of housing and force low-income residents to relocate (Wolch et al., 2014). Addressing environmental justice issues requires participatory planning and community-based strategies to address the structural changes that may be required (e.g., by improving the access of marginalized groups to green spaces and providing them with opportunities for recreation, urban agriculture, flood protection, urban heat reduction, and other ecosystem services without forcing the displacement of affected groups).

⁸ Environmental justice is a normative concept and social movement concerned with the spatial distribution of environmental goods and ills (Ernstson, 2013), as well as with the social structure and institutional context in which environmental decisions are made (Cole and Foster, 2001).

Case Study 8.7 Singapore’s Ecosystem-Based Adaptation

Lena Chan, Geoffrey W. H. Davison

National Biodiversity Centre, National Parks Board, Singapore

Keywords	Flood risks, ecosystem-based adaptation, resilience, greeneries
Population (Metropolitan Region)	5,607,300 (Department of Statistics Singapore, 2016)
Area (Metropolitan Region)	719.2 km ² (Department of Statistics Singapore, 2016)
Income per capita	US\$51,880 (World Bank, 2017)
Climate zone	Af – Tropical rainforest (Peel et al., 2007)

SUMMARY

Singapore has taken an integrated and interdisciplinary approach to urban biodiversity conservation and restoration of ecosystems by adopting both biological and engineering approaches to climate change adaptation and mitigation. Multidimensional strategies are planned and implemented to address multiple climate stressors such as temperature and sea level rise and increased water-induced hazards. Restoring terrestrial and marine biodiversity through both *in-situ* and *ex-situ* conservation work and building green infrastructure such as urban parks, wetlands, and roadside avenues have increased urban greeneries and carbon sequestration and reduced flood disaster risks. These integrated ecosystem-based adaptation and disaster risk reduction measures have made a resilient Singapore.

CASE DESCRIPTION

Singapore has taken a holistic approach to addressing climate change vulnerability and impact to its urban ecosystems. It carried out two national climate change studies incorporating vulnerability assessments that investigated physical and meteorological parameters by using statistical and/or dynamical downscaling to better understand the implications of latest Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report (AR5) climate change projections at regional and local levels⁹ (National Climate Change Secretariat [NCCS], 2012). It was followed by studying a range of downstream impacts that fed into adaptation plans based on a risk assessment exercise done across all government agency levels (NCCS, 2012).

The aim of adaptation and mitigation plans includes reducing emissions across sectors (NEA, 2013), building capabilities to adapt to the impact of climate change, and harnessing green growth opportunities, as well as forging partnerships on climate change actions. The approach assesses Singapore’s physical vulnerabilities to climate change based on a resilience framework (RF) to guide measures against potential climate change impacts. The RF ensures that appropriate adaptation measures are identified and implemented by

adopting a cyclical approach to risk appraisal and adaptation planning. The cycle is shown in Case Study 8.7 Figure 1.


CLIMATE CHANGE IMPACTS ON BIODIVERSITY

While assessing risks and planning adaptation measures, an understanding of biological and environmental assets was gained through risk identification and quantification. Biodiversity assets are understood through continuing surveys, such as site- or habitat-specific studies including the Terrestrial Sites Survey (2002–2003), Natural Areas Survey (2005–2007), and Comprehensive Marine Biodiversity Survey (2011–2015) to update information on the flora and fauna of Singapore. Regular and *ad-hoc* assessments of biodiversity and environmental assets are undertaken as part of long-term adaptation planning.

VULNERABILITY AND IMPACTS ASSESSMENT AND ADAPTATION PLANNING

The first vulnerability assessment looked at plant groups particularly vulnerable to climate change, such as figs (as keystone species for vertebrates), *dipterocarp* trees (whose bi-annual mass flowering events are keyed to the intensity and frequency of El Niño events), bryophytes (group susceptible to drought), and the effects on planted roadside trees.

Challenges have been encountered in the administrative definition and categorization of natural assets (e.g., whether each tree, each species, or each population in different areas is to be considered a separate asset) and in suggesting biological thresholds or tipping points that might be related to the various climate change parameters (rainfall, sea level, sea surface temperature, wind) in a way that facilitates risk assessment.



Case Study 8.7 Figure 1 Cyclical approach to adaptation planning.

⁹ The Strategy, developed with public and private-sector consultation, is available at <https://www.nccs.gov.sg/sites/nccs/files/NCCS-2012.pdf>



Case Study 8.7 Figure 2 *A well-planned urban ecosystem.*

Past fragmentation of Singapore's forests makes them vulnerable to future long-term changes such as increased likelihood or duration of drought and higher average temperatures. Wetlands are exposed to rainfall changes, sea level rise, or water quality changes related to warming and changes in precipitation. Sea level rise will be a challenge for mangroves, which cannot retreat inland because of competing land uses. Corals, which require sunlight, might not be able to grow upward quickly enough to keep pace with rising sea levels. In addition, a 1–2°C rise in sea water temperature will lead to coral bleaching. The strategies adopted to build up the resilience of these taxonomic groups are to conserve as broad a spectrum of species as possible and to safeguard known sources of propagules.

ECOSYSTEM-BASED ADAPTATION

To help Singapore's biodiversity withstand the potential impacts of climate change, National Parks is working with other agencies and the community to safeguard existing species, increase connectivity of various green areas across the island, and enhance the resilience of ecosystems. This includes measures to restore forest and mangrove areas through planting and through minimization of other pressures (e.g., by removing alien invasive competitors and controlling ship wakes on mangrove coasts). Singapore has a very high proportion of planted roadside trees; efforts are made to diversify plant species used, intensify planting, create more complex 3D layering, and increase connectivity between green areas.

To keep the city green, tree management and maintenance is being intensified and enhanced. National Parks manages approximately 350 parks and 3,500 kilometers of roadside greenery islandwide as part of the effort to lower ambient temperatures. Parks and greenery are not viewed as merely the passive victims of climate change, but



Case Study 8.7 Figure 3 *Innovations in urban greenery development.*

as tools for adaptation and mitigation. In addition, National Parks continues to support research that investigates the responses of coral reef communities to climate change triggers and promotes strategies that increase biodiversity resilience.

BUILDING A RESILIENT WATER SYSTEM

Water resource management is a key priority for Singapore. An increase in weather variability may bring more frequent or more severe cycles of floods and droughts threatening the reliability of the city's water supply. To ensure a sustainable water supply for Singapore's population and industry, Singapore has built a robust and diversified water supply through four "national taps": namely, local catchment water, imported water, treated recycled water (NEWater), and desalinated water. In particular, NEWater and desalinated water are not dependent on rainfall and are thus more resilient sources in times of dry weather. Regarding flood water management, efforts are made to enhance resilience against coastal erosion and inundation associated with rising sea levels coupled with short-lived, extreme meteorological events (MEWR, 2014). A risk map study was done to better identify the specific coastal areas at risk of inundation and the potential associated damage. The results will help develop long-term coastal protection strategies.

Recognizing the impact of greater weather uncertainties as well as the constraints to drainage planning posed by increasing urbanization, Singapore has revamped its drainage management approach to strengthen its flood resilience. The strategy is to optimize the management of stormwater using a holistic source-pathway-receptor approach that looks at catchment-wide ecosystem-based solutions to achieve higher drainage and flood protection standards. It covers the entire drainage system, addressing not just the pathway over which the rainwater travels. A new provision has been added to its surface-water drainage regulations, which requires developers/owners of land size 0.2 hectares or more to implement measures to slow surface runoff and reduce the peak flow of stormwater into the public drainage system by implementing on-site detention measures such as green roofs, rain gardens, and detention tanks.

CONCLUSION

Comprehensive and ecosystem-based adaptation strategies used by Singapore to enhance urban resilience are broad and interdisciplinary. These are approached from a multi-agency and multidisciplinary perspective. Additional efforts are continuously entrained in coordination with agencies and development partners under a common framework on risk, adaptation, and mitigation.

Pascual et al. (2010) argue that the institution of payment for ecosystem services (PES) is another policy area where distributive justice has critical importance. Although PES theory commonly disregards distributed justice questions, actual programs are often required to take such issues into consideration for legitimacy and stakeholder buy-in. Depending on the fairness criterion used (e.g., equal distribution, need, compensation), the outcome of PES programs are determined by an equity–efficiency interdependency analysis (Pascual et al., 2010). By including a fairness criterion of some kind, programs can offer a mechanism to more systematically include equity and justice issues in management and planning for UES (Salzman et al., 2014).

Climate change effects in coastal cities expose the complexities and challenges of developing policy to address issues of distributive justice. For example, in New York, Hurricane Sandy in 2012 devastated many coastal communities. Federal, state, and city programs determined the redevelopment path of such communities with some areas purchased for the purpose of creating new protective natural coastal buffer zones (NYS, 2013). This effort aimed at improving the adaptive capacity of the entire city to future extreme weather events. Some low-income urban residents were unable to rebuild their houses and had to relocate (Sandy Redevelopment Oversight Group, 2014). In other cases, newly required building elevations and other building reinforcement policies may mean that the individual's or community's ability to pay determines whether a family is able to rebuild its residence or instead has to relocate (Consolo et al., 2013). Planning decisions can, be complex, such as determining how best to serve residents in low-income countries where informal urban settlements are located in flood-prone areas, thus emphasizing the need to consider the broader complexity of the social, ecological, and economic linkages of the urban system.

8.8.3 Economics of Ecosystem-Based Adaptation and Green Infrastructure

Green infrastructure and other types of urban ecosystems in urban areas generate monetary and nonmonetary value through the provision of ecosystem services (Gomez-Baggethun et al., 2013; Kremer et al., 2016b) (see Chapter 7, Economics, Finance, and the Private Sector). A major advantage of green infrastructure and EbA strategies is that they offer some of the most cost-effective adaptation options available to cities (TEEB, 2011). Around the world, evidence is mounting that effective planning, design, and management of nature in urban areas can provide multiple benefits and cost-effective solutions where traditional "gray" infrastructure solutions alone have been prohibitively costly. Linking green and gray infrastructure can provide cities both cost-effectiveness and improved function.

For example, management of stormwater runoff through green infrastructure is becoming increasingly popular among cities due to the cost savings it provides by reducing the need for new gray infrastructure to reduce local flooding and sewage overflows in combined sewage systems (see Case Studies 9.4 or 14.B, Rotterdam). Green infrastructure methods such as green streets, tree plantings, and rain barrel installations are estimated to be three to six times more effective for stormwater management than further expanding gray infrastructure (Foster et al., 2011). A U.S. Environmental Protection Agency report analyzing thirteen case studies from cities such as New York, Philadelphia, Portland (Oregon), and Seattle found that although each municipality or entity used different cost and benefit matrices, in most cases green infrastructure was found to cost less than gray alternatives and to provide multiple benefits (NYC, 2010). Portland's Cornerstone project to disconnect downspouts resulted in the removal of approximately 1.5 billion gallons of runoff from

the city's combined sewer system (Foster et al., 2011), and, in Philadelphia, more than 100 green acres were constructed and 3,000 rain barrels distributed to support increased stormwater absorption. A life cycle analysis of Low Impact Development (LID) in a New York neighborhood found a strategy that included permeable pavement and street trees to be cost effective even though it only considered energy saving in downstream treatment plants; this has mirrored similar studies conducted in other cities.

Other important examples of ecosystem services include flood risk reduction by extending time lag between floods and

storm runoff and temperature regulation, ground water recharge, and air purification. Rezoning areas for green infrastructure or restricted development are cost-effective ways to address flood risks (Foster et al., 2011). Kousky et al. (2013) evaluate avoided flood damages against the cost of preventing development on flood-sensitive lots in Wisconsin and New York. Their findings highlight the importance of the spatially specific characteristics of the lot as a way to create a cost-effective flood protection plan.

UHI research shows that the loss of urban vegetation increases the energy costs of cooling (McPherson et al., 1997).

Case Study 8.8 Seattle's Thornton Creek Water Quality Channel

Nate Cormier

SvR Design Company, Seattle


Keywords	Stormwater treatment, green infrastructure, public space, ecosystem-based adaptation
Population (Metropolitan Region)	3,613,621 (U.S. Census Bureau, 2010)
Area (Metropolitan Region)	15,209 km ² (U.S. Census, Bureau, 2010)
Income per capita	US\$56,180 (World Bank, 2017)
Climate zone	Csb – Temperate, dry summer, warm summer (Peel et al., 2007)

The Thornton Creek Water Quality Channel, located at the headwaters of the South Branch of Thornton Creek, Washington, is a multi-purpose water management project providing multiple environmental and social benefits to the urban population of Seattle. This facility addresses the problem of both heavy sedimentation and polluted water flow into the natural creek in the hilly catchments of Seattle. The integrated water treatment and management plant captures runoff from the human-populated upstream watershed areas and treats it before it flows into Thornton Creek and Lake Washington. The environmentally sound water cleaning facility occupies minimal space but provides multiple spatial and environmental benefits to the local community. It has also led to the development of a new neighborhood that is emerging as a growing urban center of the city. The facility can be termed as classic example of urban green infrastructure.

INNOVATIVE AND RESILIENT DESIGN

The project uses natural drainage system revival technology simulating the natural process of water flow to clean polluted and silted water and allow the cleaned water to flow through natural percolation and seepage systems year-round. The environmentally friendly design (Case Study 8.8 Figure 1) has developed natural landscaping and public pathways giving easy access to citizens to different public facilities and private buildings located throughout the area.

This model project offers the last-available opportunity to improve the quality of stormwater runoff before it reaches the creek. The channel design diverts stormwater from the drainage pipe under the site to a series of surface swales landscaped with special soils and



Case Study 8.8 Figure 1 An innovative natural drainage design.

native plants. These ponds interrupt runoff speed, allowing water to seep into the soil and removing pollutants in the process. The channel regulates the water flow both during wet and dry weather, allowing for continuous cleaning of stormwater.

The community-driven project turned into a collective action effort that met the broad objectives of major stakeholders and fulfilled their common goals. The design has allowed development of diverse types of residential buildings, job-creating private-sector enterprises, retail shops, and rest and recreation places while preserving a natural environment. This is in contrast to what existed before – a gray and brown parking lot. The provision of public open space has been used to raise environmental awareness thus providing long-term benefits, albeit of intangible nature. The facility has attracted significant private-sector investment in terms of the residential and commercial complex. The modest US\$14.7 million that it cost to build the Thornton Creek facility is believed to have generated more than US\$200 million in the form private-sector-led investment in the city, thus catalyzing the Northgate neighborhood as a vibrant urban center of Seattle (Benfield, 2011).

ADAPTATION STRATEGY

Carved out of a former mall parking lot, the Thornton Creek Water Quality Channel provides public open space for Seattle's Northgate neighborhood while treating urban stormwater runoff from 680 acres of North Seattle. This project grew out of grassroots efforts to transform the piped Thornton Creek that ran under the parking lot to a natural water catchment system. Political leaders overcame a number of barriers that stood between developers and environmentalists by establishing a broad-based Northgate Stakeholder Group to find a way to integrate private development, public open space, and a major stormwater facility. What resulted through these collective efforts is an adaptive and resilient urban ecosystem management

project providing multiple climate change adaptation and social benefits.

Opened in 2009, this catalytic natural space provides pedestrian connectivity among a major transit hub, community services, housing, and retail outlets. There is a continuous expression of water flowing, pooling, and cascading in the channel. During and after storms, the full capability of the broad channel bottom is engaged for water-quality treatment. Overlooks and bridges allow users to enjoy the channel habitats and wildlife. Seat walls, benches, and interpretive artwork contribute to an inviting environment where visitors can linger and learn in a high-performance landscape (see Case Study 8.8 Figure 2).

The project has resulted in:

- A successful community process that balances public and private goals in support of environmentally compatible development and socioeconomic sustainability developed in a highly contested urban space

- The ability to catalyze more than US\$200 million in investment in adjacent private residential and commercial development, generating jobs and economic opportunities
- An illustration of how to transform a former mall parking lot, a common “grayfield” in many American communities, into an aesthetically and environmentally productive urban landscape.
- Water-quality treatment for runoff from 680 acres within a beautiful setting where visitors can learn about natural systems and the restoration of a historic creek.
- Increases in open space in the Northgate Urban Center by 50% to provide an oasis of native vegetation for neighbors and wildlife, thus promoting urban biodiversity.

The key lessons learned are that (1) multistakeholder processes and community-driven initiatives lead to change in developing urban resilience, and (2) both bottom-up and top-down processes are necessary, provided the city government recognizes and internalizes both in urban ecosystem-based adaptation planning and implementation.



Case Study 8.8 Figure 2 *Thornton Creek in Seattle: An example of human-developed biodiversity and ecosystem.*

Significant savings can accrue due to the reduction of power generation through the implementation of green infrastructure (ACCCRN, 2015; Rosenzweig et al., 2009). For example, one study in Los Angeles showed that increasing pavement reflectivity by 10–35% could produce a 0.8°C decrease in UHI temperature and an estimated savings of US\$90 million per year from lower energy use and reduced ozone levels (Foster et al., 2011).

8.8.4 Payment for and Valuation of Ecosystem Services

Many cities are developing programs for valuation of and payment for ecosystem services. However, PES programs are

not often decided based on proper valuation and often fail to address the issue of social equity; in some cases, they exacerbate poverty and equity by raising prices or introducing a fee on previously low-priced or free services (Pascual et al., 2009).

Common valuation methods include preference-based approaches and biophysical approaches (Sukhdev et al., 2010). *Preference-based approaches* include all monetary and nonmonetary societal value settings, and *biophysical approaches* include assessments that are grounded in the processes, flows, and structures of the ecosystem (Sukhdev et al., 2010; Gomez-Baggethun et al., 2013). An important characteristic of urban green infrastructure is that it generates multiple benefits and different types

of values (Kremer et al., 2016b). One of the challenges in the evaluation of the cost-effectiveness of green infrastructure is accounting for societal and cultural benefits and values that are not easily quantifiable in monetary terms. For example, a study of flood protection strategies in the Netherlands found engineering methods to be most cost-effective when not considering nonmonetary benefits, but when those were included (e.g., social and cultural values), green infrastructure became more competitive. Such integration is at the forefront of current UES research (Haase et al., 2014).

8.8.5 Economic Valuation Tools

Because of a growing effort to support the integration of green infrastructure into the urban landscape, software tools are becoming increasingly available to urban planners and decision-makers for the evaluation of certain ecosystem services and benefits. For example, i-Tree¹⁰ is a suite of software tools built by the U.S. Department of Agriculture Forest Service that allows the quantification of ecosystem services benefits from urban trees; the Green Values Calculator¹¹ is a tool for comparing performance, costs, and benefits of green infrastructure practices; and InVEST¹² is a suite of software models for the assessment of ecosystem services values and tradeoffs (Nowak et al., 2013). Such tools enable the valuation of UES and support the integration of green infrastructure into urban planning. However, major gaps remain in the capacity to value urban green infrastructure and the ecosystem services it provides, including public participation in the valuation process, the integration of monetary and nonmonetary values through multicriteria analysis and other methods, scale- and thresholds-dependent values, and bridging supply and demand for the purpose of valuation (Gómez-Baggethun et al., 2013; Haase et al., 2014). Additionally, costs of EbA and nature-based solutions for climate mitigation and adaptation often have to be estimated, especially with respect to future costs, since adaptation is a long-term process. In most cases, obtaining reliable cost data will continue to be a challenge requiring several sources of evidence ranging from project case studies to national-level assessments.

8.8.6 Combining Adaptation and Mitigation in Climate Resilience Strategies

Although adaptation is necessary to minimize the unavoidable impacts of climate-induced risks and hazards, mitigation is needed to reduce urban GHG emissions and their impacts in the short- and long-term. An integrated strategy that combines all types of adaptation and resilience building measures together with mitigation strategies will have the highest level of co-benefits for human well-being (Satterthwaite et al., 2008; Karki et al., 2011) (see Chapter 4, Mitigation and Adaptation). Risks and vulnerabilities are shaped by local environmental conditions,

site characteristics, natural resource availabilities, and environmental hazards (IPCC, 2007; Satterthwaite et al., 2008). Urban adaptation aims at reducing vulnerabilities and enhancing the resiliency of systems, agents, and institutions, and it needs to be planned by taking a holistic view of the broader urban landscape since urban areas depend on surrounding peri-urban and rural areas for ecosystem services (Tyler and Moench, 2012).

Strategies for urban ecosystem adaptation and mitigation need to recognize that climate change may undermine the ability of contiguous urban and peri-urban social ecological systems to provide critical ecosystem services (Satterthwaite et al., 2008). Therefore, urban adaptation and mitigation planning should ensure the sustained flow of provisioning (e.g., food, water) and regulating (e.g., clean air) ecosystem services to urban communities (Locatelli et al., 2010; McPhearson et al., 2015). In many parts of the world, the relationships among urban ecosystems, adaptation, mitigation, and livelihoods are changing in fundamental ways as urban economic systems diversify across the urban–peri-urban spectrum, thus creating mixed or interlinked economic and environmental systems. Understanding these changes and their implications on the vulnerability of urban populations and ecosystems is essential to developing integrated adaptation and mitigation strategies for cities.

For example, Singapore has taken steps to restore its biodiversity and enhance UES (see Case Study 8.7, Singapore). The city has increased green cover from 35.7% to 46.5% in 20 years and also has set aside approximately 10% of its total land for green infrastructure (Lye, 2010) to provide increased climate change mitigation in the city in addition to improving ecosystem services that support adaptation. Similarly Seattle, Edmonton, Stockholm, Copenhagen, and many other cities have restored or created new urban ecosystems that ensure a more sustainable flow of ecosystem goods and services to the city-dwellers now and in the future (Zandersen et al., 2014).

8.8.7 Biodiversity Governance for Human Well-Being in Cities

In many parts of the world, the relationship between urban ecosystems and overall urban development is changing in fundamental ways (Tzoulas et al., 2007). Understanding these changes and their implications is necessary for holistic sustainable urban development. Specifically, over the coming decades, two interacting forces will influence urban economic and ecological systems especially in developing countries: (1) intensifying processes of technological and economic globalization that are already increasing pressures on urban/peri-urban ecosystems through shifting patterns of dependency; and (2) multiple environmental stress at all levels – from local to regional – mainly due to the impacts of climate change. These changes will likely undermine the ability of complex urban ecosystems to provide

10 <https://www.itreetools.org>

11 <http://greenvalues.cnt.org/national/calculator.php>

12 <http://www.naturalcapitalproject.org/InVEST.html>




Figure 8.8 Overview of global governance arrangements for urban biodiversity and ecosystem services.

Source: UN-Habitat, 2012

critical services – water, energy, food, clean air, and healthy and livable habitats – to their population, thus underlining the critical importance of urban planning, policy, and governance to safeguard urban and peri-urban biodiversity and ecosystem services (see Chapter 16, Governance and Policy).

The decisions and deliberations of the UN Convention on Biological Diversity (CBD, 2009), as well as many others (ICLEI-Africa, 2013; UNEP, 2009; IUCN, 2009), have created an emerging global effort to enhance urban ecosystem governance structures by capturing the nexus between urban biodiversity and climate change. These efforts have called for biodiversity vulnerability assessments and research on links between biodiversity loss and urbanization (Wilkinson et al., 2010, 2013; Schewenius et al., 2014). Figure 8.8 illustrates the current international governance landscape for urban biodiversity and ecosystem services, although it is constantly evolving. Progress remains at a formative stage as stakeholders and urban actors struggle to fully understand their respective roles and establish coordination mechanisms to exploit the latent potential

of biodiversity and ecosystem services in cities for climate change adaptation and mitigation.

Increasing a city’s capacity to meet growing challenges can depend on the development of a holistic governance approach in which the city is understood as a dynamically interacting social-ecological system (Frantzeskaki and Tilie, 2014) (see Chapter 1, Pathways to Urban Transformation and Chapter 16, Governance and Policy). Increased linkages among strategies, projects, and actors (Meyer et al., 2012), including the active involvement of local citizens, is important for identifying needs, challenges, and design policies and efficiently implementing them (Ward et al., 2013; Wardekker et al., 2010). Creating strong links between formal governance and informal, on-the-ground participants and managers is crucial to forming holistic governance with greater potential for successful urban ecosystem management outcomes. However, informal participation and management is seldom translated into formal governance in urban settings (Colding, 2013). The Thornton Creek Water Quality initiative of Seattle is an example that shows how a local

community-driven project supported by city government is generating multiple benefits and synergies (see Case Study 8.8).

Governing ecosystem processes requires coordination across different levels of policy, legislation, and jurisdictional authorities. Urban ecosystems and biodiversity benefits often transcend administrative boundaries, thus necessitating collaboration among national, regional, and local-level agencies (Ernstson et al., 2010; McPhearson et al., 2014). The multiscale and multi-sectoral relationships that impact urban biodiversity and ecosystem services create urban governance and policy challenges because decisions by one branch and one level of fragmented urban and national government structures can create long-term implications for the entire urban ecosystem landscape (Asikainen and Jokinen, 2009; Ernstson et al., 2010; Borgström et al., 2006). Apart from this scale mismatch issue, there is also a functional mismatch between ecosystems and the institutions managing them (Cumming et al., 2006), because different decision-makers are operating within and beyond the city and urban landscape boundaries. However, if different units of city and peri-urban governments worked in tandem, a number of synergies in the governance of urban biodiversity and ecosystem services is possible (Raudsepp-Hearne et al., 2010). This requires functioning and dynamic science–policy linkages at regional scales, challenging the current structure of governance frameworks, practices, and institutions.

Despite new research initiatives and science–policy platforms, significant challenges remain in equitably managing biodiversity and ecosystem services in urban and peri-urban areas for the mutual benefit of humans and other species (Schewenius et al., 2014). Rapid urbanization is occurring in places that face some of the most severe challenges to public health and urban biodiversity conservation. Additionally, these same urban systems are where systems of formal government and planning tend to be weak and the capacity to influence policy inadequate (Wilkinson et al., 2013).

Effective city governance will play a key role in determining the future of biodiversity across the world, not least because cities are rapidly expanding into the world's biodiversity hotspots (see Figure 8.5). Significant urban ecosystem policy changes will need to accompany or even precede effective governance practices in order to direct future urban growth so that biodiversity and the ecosystem services it provides are safeguarded (Seto et al., 2012; Wilkinson et al., 2013). Ecosystem protection in cities will rely on increasing efforts by parks and natural area managers to focus on those management outcomes that seek to maximize ecosystem functioning for services – in many places an abrupt shift from existing or past management goals. Supporting a diversity of governance systems, from official regulations to informal governance systems (e.g., local governance of urban allotment gardens) can provide a multilayered protection system and strengthen support through multiple stakeholders (Schewenius et al., 2014). Additionally, flexible policies and regulations will be needed to accommodate unanticipated climate changes and ecological responses.

8.9 Science–Policy Linkages

Adaptive and resilient urban ecosystem conservation requires policies that are based on synthesized and relevant knowledge systems, local evidence, and multilevel, multidisciplinary, and multi-stakeholder consultations and inputs (IUCN, 2009; Van Kerkhoff and Lebel, 2006) (see Chapter 16, Governance and Policy). Three important challenges inhibit the use of climate and conservation science in managing and planning urban ecosystems: (1) research capacity and activities are often scattered so that policy-influencing efforts are uneven in distribution and quality across sectors and regions (Haase et al., 2014); (2) data and information availability and usability are limited due to knowledge gaps and scale-appropriate specificity (e.g., in Sweden, local planners and decision-makers found it difficult to implement national biodiversity strategies because they were too general and abstract); and (3) policy-makers and other decision-makers often have limited capacity to access, interpret, and act on research information on technical subjects such as biodiversity and climate change, particularly where results are complex and reflect inherent uncertainties at multiple scales (Amin, 2007). Urban policy-makers therefore face significant challenges when seeking to increase the resilience of communities and the built environment to the effects of climate change (Frantzeskaki et al., 2016).

An integrated and coordinated urban planning, design, and implementation policy that considers biodiversity and ecosystem services should address multiple co-benefits from human health improvement, climate change adaptation, biodiversity conservation, and disaster management (see Chapter 5, Urban Planning and Design). For example, increasing green space, tree cover, and water bodies in urban areas, in addition to moderating UHI effects, will also sequester carbon, control pollution such as aerosol dusts, regulate hydrological processes, and influence regional climate (Hamstead et al., 2015; Larondelle et al., 2014). Policy actions that take advantage of the complex concepts of multifunctionality, synergies, and tradeoffs require science- and evidence-based policy processes that integrate ecosystem-based approaches in governance areas such as disaster management, community actions, and linking adaptation with sustainable development goals and practices (Elmqvist, 2013). Such processes involve key urban actors (including politicians) and civil societies in urban conservation activities that can translate research-based information and knowledge into use by policy-makers and other decision-makers (Elmqvist et al., 2013; Mincy et al., 2013; OECD/CDRF, 2009).

Co-production of knowledge, where the users of the knowledge are involved from the beginning in the research and review process, is another key component of successful science-based policy-making. The Asian Cities Climate Change Research Network (ACCCRN) has concluded that urban process needs to be based on multiple-stakeholder engagement and iterative shared-learning dialogue that can bring a broad range of perspectives to city managers. Urban policies are rarely completely objective or neutral. Shared learning processes and co-production of knowledge can help raise awareness and empower stakeholders with new and consolidated

Box 8.6 WWF's Earth Hour City Challenge

The World Wildlife Fund (WWF) supports a vision of the world where people and nature thrive. In an increasingly urbanizing world, achieving this vision means working together with cities to make them livable and sustainable. WWF is the world's largest conservation organization, working not only on wildlife protection but also on food, oceans, forests, water, and climate change. WWF is bringing together its network of experts on renewable energy, public engagement, nature-based adaptation, and many other disciplines to address the issues cities are facing in the 21st century, in particular the threat of climate change and its associated hazards.

WWF's signature program for cities is the Earth Hour City Challenge (EHCC). EHCC was created in 2011 to mobilize action and support from cities in the global transition toward a sustainable future. It has since grown to encompass cities in twenty countries around the globe. Last year, 166 participating cities reported their climate data, commitments, and a total of 2,287 mitigation actions on the carbon Climate Registry (cCR) for review by an esteemed jury of experts. The jury, comprising high-level representatives from key city networks, development banks, institutions, universities, and enterprises, evaluate the participating cities' goals and strategies. Every year, one city from each participating country is awarded the title National Earth Hour Capital. From among these inspiring finalists, the jury then selects one Global Earth Hour Capital. WWF offices in twenty countries support cities on EHCC communications and low-carbon project implementation.

One key objective of the EHCC has been to gather a critical mass of city reporting on their climate commitments;

and climate actions in order to raise the awareness of decision-makers involved in global climate negotiations and increase aspirations and actions at the national level.

The We Love Cities campaign profiles finalists and spurs interaction between cities and their citizens through social media. Public engagement and raising awareness around the positive stories on local climate action are key components of the program. We Love Cities invites citizens from around the world to express their love through votes, tweets, and Instagram pictures and by submitting suggestions on how their cities can be more sustainable. These suggested improvements are shared among all the participating cities. More than 300,000 people who truly love their cities and want to see them become more sustainable have engaged in this campaign.

WWF works closely with the ICLEI–Local Governments for Sustainability to run the EHCC as well as many country-level programs that extend technical and communications support to cities around the world. In addition to ICLEI, WWF is partnering with other leaders to address climate change including C40 Cities Climate Leadership Group, Compact of Mayors, Rockefeller 100 Resilient Cities, U.S. Agency for International Development.

Technical guidance and original research from WWF are also available to support cities including the Green Recovery and Reconstruction Training Toolkit; Green Flood Risk Management Guidelines; Urban Solutions for a Living Planet; Measuring Up 2015; Financing the Transition: Sustainable Infrastructure in Cities; and Reinventing the City.

knowledge (Sutcliffe and Court, 2005; Institute for Social and Environmental Transition [ISET], 2010).

Strategies to effectively link science with policy and action need to (1) involve key actors (local residents, planners, designers, managers, policy-makers, NGOs) in the process of identifying problems and the actions they can take (i.e., shared learning and knowledge coproduction), (2) produce grounded evidence where action can be used to respond to ecosystem changes that are relevant to members of communities and key sectoral decision-makers, (3) effectively communicate evidence to an array of end users so that they understand and can act on it (translation of research results into use depends critically on how they are communicated via direct experience, accessible products, and, for academic and policy global audiences, peer-reviewed articles), and (4) design research outputs to respond to the types of information different types of actors need and can relate to (e.g., cost-benefit analyses and regulatory regimes for government and multilateral investors, new business opportunities for the private sector, equity concerns for the community groups, and examples of tangible solutions to common climate vulnerabilities that individuals and households face). These approaches can help to build incremental science–policy linkages that support efforts to transition cities toward sustainability and resilience.

8.10 Knowledge Gaps and Areas for Further Research

Sustainable generation and management of urban biodiversity and ecosystem services in the face of the challenges posed by climate change, population growth, poverty, and environmental degradation requires adaptive human and institutional capacity that can enhance resilience, human well-being, and conservation (TEEB, 2010; Elmqvist et al., 2013; RUAF, 2014). However, a common problem that urban policy-makers and city managers face when dealing with climate change is bridging significant knowledge gaps. This is especially challenging in the context of climate change effects on cities and urban areas (Elmqvist et al., 2013). The release of the IPCC AR5 report made headway in bridging the knowledge gap at the global level (IPCC, 2014), but for many urban areas, especially in developing countries, data and knowledge gaps remain a problem at both local and regional levels. Enhancing urban biodiversity and ecosystem services while tackling climate change and a host of social issues in cities requires a continuous flow of knowledge-based solutions (see ARC3.2 Annex 1, UCCRN Regional Hubs). Missing empirical evidence and practical ecological knowledge on urban biodiversity and ecosystems management often prevents city managers

from recognizing the value of ecosystems for the development of more climate-resilient urban systems.

Additionally, significant knowledge gaps remain in understanding the current status of biodiversity in cities. Despite growing databases and new global analyses of urban biodiversity and ecosystem services (Elmqvist et al., 2013; Gomez-Baggethun et al., 2013; Aronson et al., 2014), most cities, especially in low- and middle-income countries do not have adequate data on the status and extent of biodiversity and urban ecosystem resources. Leveraging UES for climate resilience is hampered by this lack of data, with multiple global and local agencies and institutions calling for national, regional, and local biodiversity and ecosystem assessments.

Producing tools and guidelines on how to effectively manage and govern urban ecosystems so that critical services are available to local populations remains an area in need of additional research and practice-based expertise (Schewenius et al., 2014). Benefits of biodiversity and ecosystems are not equally distributed in urban areas. Often, poor communities or housing for them (e.g., Cape Flats) are blamed for biodiversity loss and habitat fragmentation in spite of their low per capita impact or having been pushed to the most marginal and fragile sites (Ernstson et al., 2010a). Improving equitable distribution and access to ecosystem services, whether it is for shade relief from urban heat waves or protection from climate-driven extreme events such as flooding in coastal cities, depends on increasing equality and reducing mismatches between ecosystem services supply and social demand for these services (McPhearson et al., 2014; Salzman et al., 2014).

Although cities and urbanized regions depend on biodiversity in ecosystems to sustain human health and well-being (TEEB, 2011), this relationship is not well understood for all ecosystem services, and the connection between biodiversity and human livelihoods has yet to be widely incorporated in urban policy and planning (Hansen et al., 2015; McPhearson et al., 2014; McPhearson et al., 2016). We also still know little about how biodiversity, ecosystem function, and ecosystem services are related in urban environments. Empirical and theoretical research on the relationships among biodiversity (including native and non-native species), ecosystem function, and ecosystem services is critical for developing design standards for climate-resilient green infrastructure.

8.11 Recommendations for Policy-Makers

The growing impacts of climate change and climate variability on interconnected human–environmental urban systems are increasing the vulnerability of both human and ecosystems in cities. Cities are particularly at risk. Ecosystems in urban contexts underpin the security of public health, water, food, industrial activities, biodiversity conservation, energy, and transport, as well as recreation and tourism sectors. Effective management of urban ecosystems using multisector and multiscale approaches

will be key in the pursuit of climate-resilient, sustainable urban development.

Adaptive management of ecosystems at landscape or watershed scales involving all stakeholders across municipal boundaries is critical to safeguarding ecological resources for climate adaptation and mitigation. Investing in green infrastructure and EbA is particularly relevant for cities and urbanized regions because they can integrate climate change adaptation and disaster risk reduction, providing cost-effective nature-based solutions for addressing climate change in cities (UNEP, 2012; Munroe et al., 2010). Investment in green infrastructure and EbA can generate multiple co-benefits for human well-being by mainstreaming climate and environmental considerations across urban systems and encouraging the sustainable management of ecological resources to improve the resiliency of inhabitants, built environments, and urban infrastructure. These approaches have the potential to mainstream environmental and climate change information into urban planning, decision-making in urban design, and management and implementation processes. Research in urban systems is making clear the cost-effective, widely beneficial impacts of investing in biodiversity and urban ecosystems for climate adaptation. We suggest the following policy-relevant recommendations:

1. Invest in ecosystem-based adaptation and green infrastructure planning as critical components of climate adaptation strategies and urban development, as well as for improved health, disaster risk reduction, and sustainable development.
2. Incorporate the monetary and nonmonetary values of biodiversity and ecosystem services into cost-benefit analyses for climate adaptation and urban development and develop innovative means of financing (e.g., public–private partnerships) for urban ecosystem and biodiversity protection, restoration, and enhancement.
3. Utilize a systems approach to ecosystem-based climate adaptation, explicitly recognizing the social-ecological relationships that co-produce ecosystem services and drive ecological dynamics in urban systems.
4. Plan and manage for a sustained supply of critical urban and peri-urban ecosystem services over longer-term time horizons (20, 50, 100 years).
5. Strengthen urban–peri-urban–rural linkages through integrated and multidisciplinary urban and regional ecosystem planning and management and involve local communities and diverse stakeholders to reduce the vulnerability of urban poor and minorities.
6. Launch collaborative, cross-boundary, and co-designed urban biodiversity and ecosystem research and advocacy programs to inform policies and planning and further develop nature-based solutions toward more resilient, livable, and sustainable urban futures.

8.12 Conclusions

Urban areas all over the globe, especially in developing countries, are growing rapidly in both population and area and are putting pressure on urban biodiversity and ecosystems to support

livability, sustainability, and climate resilience. Climate change and its impacts on cities amplify the effects of urban stressors for ecosystems. Urban biodiversity and ecosystems will need to be safeguarded and enhanced to support climate mitigation and adaptation efforts and deliver critical, nature-based co-benefits for human well-being in cities. Urban ecosystems can help offset the worst impacts of climate change, including reducing the impact of extreme events by regulating hydrology, moderating local temperature, and providing critical ecosystem services. City leaders need to recognize the interdependence of the city with peri-urban and rural surroundings and continue to broaden their planning horizon to regional levels to account for the fact that species, ecosystems, and people cross municipal boundaries and so must planning, management, and governance.

Urban and peri-urban ecosystems provide critical natural capital for climate change adaptation in cities and urban regions. Ecological spaces in cities, including all forms of green infrastructure, provide important ecosystem services such as UHI reduction, coastal flood protection, and stormwater management. Urban ecosystems are already and will continue to be affected by climate change. Cities should utilize, protect, and restore these ecosystems when seeking to improve urban resilience to the effects of climate change. City planners, managers, and decision-makers can utilize nature-based solutions to design and implement climate adaptation and mitigation strategies in combination with more traditional built infrastructure solutions. Investing in natural capital is a cost-effective strategy that also generates multiple co-benefits that enhance human well-being. In this way, urban ecosystems simultaneously provide means for improving urban resilience, livability, equity, and sustainability.

Building climate-resilient urban communities entails a socio-ecological framework as opposed to socio-technological approaches (Berkes and Folke, 1998) that can reconnect cities to the biosphere (Andersson et al., 2014). Investing in urban ecosystems for climate adaptation and mitigation makes good sense because it is cost-effective and provides numerous co-benefits that can improve equity and livability in cities. Mounting evidence of the benefits of urban ecosystems as nature-based solutions calls for strengthening climate resiliency by investing in good governance, flexible institutions, and collaborative programs. We find through this review that urban biodiversity and ecosystem services are critical to the development of climate-resilient cities.

Annex 8.1 Stakeholder Engagement

To better gauge stakeholder understanding of urban ecosystems and biodiversity for climate change adaptation and mitigation, a two-pronged approach was used to reach a diverse array of groups. Chapter authors met informally with stakeholders at workshops and meetings in Berlin, New York, Rotterdam, Stockholm, and Paris, engaging with a multidisciplinary, global group of actors who contributed broad perspectives on urban climate change and development issues. Despite these engagements being held in the United States and Europe, the stakeholders

involved were geographically, gender, and ethnically diverse, capturing views of managers, designers, citizens, planners, and policy-makers and other decision-makers. Additionally, an electronic survey was conducted to gather the views of a wider community for more formalized engagement with stakeholders (see Annex 8.2). The goal of the survey was to better understand how a broad range of stakeholders perceives urban ecosystems, their value, and their role in reducing climate change impacts and improving the resilience of cities.

Annex 8.2 Urban Ecosystem and Biodiversity Stakeholder Engagement Survey

Authored by Helen Santiago Fink, Quynn Nguyen, and Chapter Lead Authors

The stakeholder survey period was October 20–November 21, 2014, during which sixty-two responses were collected and then analyzed.

This basic survey instrument (Exhibit 8.A) was designed as part of the research for this chapter to solicit information from two key stakeholders groups (see Exhibit 8.B for Stakeholder List): (1) Urban professionals/practitioners or those entities involved in shaping or influencing the physical urban space, including planners, architects, engineers, political/regulatory decision-makers, real estate and construction industry professionals, environmental NGOs, and others; and (2) Urban end-users, which includes everyone who uses and benefits from the urban environment, from the general public to households; visitors; business enterprises; social, service, and learning facilities; and many others. The survey was developed by a subgroup of the chapter authors and reviewed by external reviewers. The twenty survey questions were structured in four parts: (1) Profile of the anonymous responder, (2) Role/value of urban ecosystem and biodiversity, (3) Relationship of ecosystems services to climate change, and (4) Socioeconomic and policy measures to support urban ecosystems services.

The survey received responses from many regions of the world, including Africa, South America, Asia, and the United States and Europe (with the largest representation). In respect to responder profiles, statistics indicated that 90% were urban dwellers; 40% were government employees; and 60% held a master's degree. Awareness of the term "urban ecosystem services" (UES) was indicated by 59% of the responders, with 28% never having heard of the term before this survey and 13% somewhat aware. The role of UES was seen by 80% of responders as valuable for aesthetics, recreation, health, pollution control, and climate mitigation and adaptation; however, climate change recorded the lowest (9%) among them all. Rural areas and wealthy populations were seen to benefit more, despite 57% of all responders acknowledging the benefits of UES for all groups. When asked to rate the value of UES among sixteen potential attributes, air quality (80%) and a healthy life (78%) followed by water quality (66%) received the highest responses. Physical,

Annex Box 8.1 Stakeholder Survey

Chapter authors pursued a two-pronged approach to engage a multidisciplinary group of stakeholders to contribute broader perspectives on the chapter's themes: (1) Consultations at relevant international conferences and workshops and (2) A detailed online survey. Although the survey had limitations in sample size ($n = 62$), it offers insights into the views of an international audience and suggests key points for broader stakeholder engagement. Key findings included:

- Stakeholders are strongly in support of protecting and enhancing UES for climate action, human well-being, and general quality of life. Despite 27% of survey respondents reporting that they had not heard of the term “urban ecosystem services,” survey results suggest wide public support for investing in urban ecosystems for climate adaptation and mitigation.
- The role of urban ecosystems services was seen by 80% of survey responders as valuable for a multitude of issues including aesthetics, recreation, health, and pollution control; however, climate change was among the lowest (9%) reported benefit.
- Stakeholders’ “strong concern for climate change” and high agreement on “associated benefits of ecosystem services” calls attention to the value of increasing awareness to better communicate the multiple benefits of urban ecosystems to society including as ecosystem-based climate adaptation and mitigation.
- Survey results indicated a favorable “willingness to pay” for ecosystem services that provide climate adaptation and mitigation.
- Research is needed to fully understand how to positively encourage “human/personal attachment to the natural environment” (acknowledged by 89% of survey responders). The goal of the stronger attachment is to promote environmental stewardship and the development of stronger policy actions and fiscal instruments to advance climate decision-making and investment for natural capital and nature-based solutions.


psychological, and spiritual well-being, as well as recreation/leisure, were rated high by approximately 53% of responders. Climatic benefits such as carbon sequestration, temperature reduction, and lessened impacts of extreme weather events (e.g., landslide prevention) received 50%. UES ratings were 21% for cultural and sport activities, and education received 48% with pollination of crops being an outlier with only one response.

Because of the small sample size of sixty-two responders, the chapter authors are cognizant of the limitations to generalizing the findings of the survey’s results. It is unrealistic to correlate the results to the wider population. Nevertheless, the survey highlights some key points for further investigation on the developing role of UES in the climate change agenda. Engagement with practitioners and decision-makers (e.g., city managers, administrators, policy-makers) at multiple levels, including with active end-users of ecological infrastructure (e.g., urban naturalists, conservationists, researchers, non-profits, NGOs, governments, social institutions, museums, community groups, and citizenry) could benefit from increased social-learning models promoting environmental education as an opportunity for increased stakeholder engagement. Communicating the critically important role that the natural environment and biodiversity play in both climate adaptation and mitigation, as well as in their nexus, is a cornerstone to elevating the climate and sustainable cities dialogue and practical action in urban areas.

In a number of separate questions, the relationship between UES and climate change was highly correlated, with 82% acknowledging a connection (see Annex Figure 8.A); similarly, 72% of responders indicated being “very concerned” about climate change. UES were perceived as important to help or be “able to protect” health (93%), water and sewage (82%), and property

values (80%), while both food supply and employment recorded a lower response (65%). There was general willingness to support UES for climate change action through economic and financial measures by around 65% of respondents. Combinations of fiscal instruments were favored by almost half of responders, yet when viewed individually, specific measures such as a carbon tax (46%), general government budget (51%), and penalty for polluters (43%) rated among the highest (see Annex Figure 8.B). The level of support was strongly recorded on a personal basis, with 68% of responders willing to volunteer and participate in a planning process for urban ecosystems for environmental protection. Regulatory encumbrances on land use were also overwhelmingly supported (62%) in order to provide more green space in cities, including restrictions on responders’ private property.

Overall, the survey suggests strong support for protecting and enhancing UES for climate action, with strong co-benefits for human well-being. The result is encouraging, given the fact that 27% of responders had not heard of the term “urban ecosystem services” before and thus indicating potentially wider multistakeholder support. The high rating for “strong concern for climate change” and a majority agreeing on reaping “associated benefits of ecosystem services” calls attention to the need for increased awareness in building efforts to educate the public on the multiplicity of benefits provided by urban ecosystems to society. The survey result indicating a general “willingness to pay” for UES to contribute to both climate mitigation and adaptation, and this suggests increasing opportunities to incorporate EbA and green infrastructure development (among other measures) into local and national urban policies and practices. Research is needed to understand how to positively exploit the strong response of a “human/personal attachment to the natural environment” (acknowledged by 89% of survey responders) toward the



Annex Figures 8.A and 8.B Results of stakeholder engagement survey: A. Demonstrates broad understanding of relationship between urban nature and climate change; B. Shows possible climate change risk reduction programs and how they are prioritized among stakeholder respondents. Highest are government budget, carbon tax, and penalty fee for polluters.

development of stronger policy actions, integrated planning, and fiscal instruments to advance climate change decision-making and investment in building climate-resilient cities.

Exhibit 8.A Urban Ecosystem and Biodiversity Stakeholder Engagement Survey

- Where do you live? (Choose one only)
 - Urban area (city) Name and Country: _____
 - Non-urban areas (suburban and rural areas) – Name and Country: _____
- What is your employment? (Choose one only)
 - Government employee
 - Private sector employee
 - Self-employed
 - Civil society/NGO/nonprofit
 - Development partner (donor) agencies
 - Others
- Which best describes your household status?
 - Single
 - Married with no children
 - Married with children
 - Other
- What is your highest level of education?
 - Post/doctoral degree
 - Master degree
 - Bachelor degree
 - High school degree (12 years of education)
 - Less than high school degree
- What in your view is the role of Ecosystems (or Nature such as trees, parks, gardens, animals, lakes, rivers, wetlands, green spaces) in a city?
 - Aesthetic value

- Recreation
- Pollution control
- Climate change adaptation and mitigation
- Others
- All

- In your view which section or who in society benefits the most from Ecosystem(s) in your community/city?
 - Rich class
 - Middle class
 - Poor class
 - Other
 - Everyone
- Which sector or community benefits most in the world from Ecosystems?
 - Urban populations
 - Rural populations
 - Global population
 - Governments
 - Businesses
- Select the importance of each of the benefits of Ecosystems/Nature in a city (rating from 1 to 5 [highest]):

(a)	Recreation and leisure	1	2	3	4	5
(b)	Sports activities	1	2	3	4	5
(c)	Cultural activities	1	2	3	4	5
(d)	Healthy life	1	2	3	4	5
(e)	Water quality	1	2	3	4	5
(f)	Air quality	1	2	3	4	5
(g)	Physical well-being	1	2	3	4	5
(h)	Psychological well-being	1	2	3	4	5
(i)	Spiritual well-being	1	2	3	4	5

(j)	Urban temperature	1	2	3	4	5
(k)	Flood prevention	1	2	3	4	5
(l)	Landslide prevention	1	2	3	4	5
(m)	Food production	1	2	3	4	5
(n)	Carbon sequestration	1	2	3	4	5
(o)	Pollination of crops	1	2	3	4	5
(p)	Others _____	1	2	3	4	5

(c)	Water and sewage	Yes	Somewhat	No
(d)	Public transportation	Yes	Somewhat	No
(e)	Employment/jobs	Yes	Somewhat	No

9. Have you heard about *Urban Ecosystem Services* before this Survey?

- (a) Yes
- (b) Somewhat
- (c) No

10. How concerned are you about Climate Change?

- (a) Very concerned
- (b) Somewhat concerned
- (c) Not concerned

11. Is there a connection between Urban Ecosystems (Nature in a city) and Climate Change?

- (a) Yes
- (b) Maybe
- (c) No

12. Can Urban Ecosystems HELP with or PROTECT the following?

12.1 PEOPLE

(a)	Health	Yes	Somewhat	No
(b)	Injuries/risks (on personal level)	Yes	Somewhat	No
(c)	Security (on community/city level)	Yes	Somewhat	No
(d)	Food supply	Yes	Somewhat	No
(e)	Water supply	Yes	Somewhat	No

12.2 ASSETS

(a)	Property values	Yes	Somewhat	No
(b)	Business losses	Yes	Somewhat	No
(c)	Foreign/domestic Investment	Yes	Somewhat	No

12.3 INFRASTRUCTURE

(a)	Roads	Yes	Somewhat	No
(b)	Electric power	Yes	Somewhat	No

13. Can expanding urban green areas in the city help prevent global warming and/or reduce greenhouse gas emissions (GHGs)?

- (a) Yes
- (b) Maybe
- (c) No

14. Can Urban Ecosystems in the city offer a better quality of life?

- (a) Yes
- (b) Maybe
- (c) No

15. Would you be willing to pay for preserving and/or expanding Urban Ecosystems to help with Climate Change risks?

- (a) Yes
- (b) Somewhat
- (c) No

16. How should a Climate Change risk reduction program through urban ecosystem improvement be funded?

- (a) Carbon tax
- (b) Penalty fee for pollution/carbon emissions
- (c) Donations
- (d) Government budget
- (e) Public-private partnerships

17. Would you be willing to participate in the process of Urban Ecosystem planning and environmental protection/conservation on a personal/volunteer level?

- (a) Yes
- (b) Maybe
- (c) No

18. Would you be willing to have more regulations/laws that would require more green spaces in public and private areas?

- (a) Yes
- (b) Maybe
- (c) No

19. Would you be willing to have a restriction(s) on your property/land in order to have a greener city/community?

- (a) Yes
- (b) Maybe
- (c) No

20. Do you feel a personal attachment to the natural environment?

- (a) Yes
- (b) No
- (c) Somewhat

Exhibit 8.B Urban Ecosystems and Biodiversity Stakeholders List

Institutional (representatives of):

Public:

1. *City (Executive)*
 - a. Climate Change – Adaptation/Mitigation
 - b. Urban Planning
 - c. Environment
 - d. Housing
 - e. Health
 - f. Education
 - g. Transportation
 - h. Urban Conservation (Infrastructure)
 - i. Parks and Gardens (Green Areas)
2. *Metropolitan/State (Executive)*
 - a. Climate Change – Adaptation/Mitigation
 - b. Environment
 - c. Economy
 - d. Transportation
 - e. Health
 - f. Education
 - g. Housing
 - h. Infrastructure
3. *City and State Councils (Lawmakers)*
Representatives of related areas (as above)

Private Economic Sectors Associations

- a. Real Estate
- b. Infrastructure
- c. Industry
- d. Commerce
- e. Transportation
- f. Tourism

Private Civil Society Associations

- a. Residents
- b. Arts and Culture, Education, Social and Political Groups, and Environmental NGOs
- c. Professionals Urban Planners, Architects, Landscape Architects, Engineers, Foresters, Agronomists, etc.
- d. Informal active organized groups

Individuals (users)

- a. Parks and trails
- b. Public transportation
- c. Pedestrians in busy streets
- d. Private transportation (car drivers)

Chapter 8 Urban Ecosystems and Biodiversity

References

- Agencia Catalana de l'Aigua (ACA). (2014) Accessed April 17, 2015: http://acaweb.gencat.cat/aca/appmanager/aca/aca?_nfpb=true&_pageLabel=P51200158891423828204014
- Ahern, J. (2007). Green infrastructure for cities: The spatial dimension. In Novotny, W, and Brown, P. (eds.), *Cities of the Future: Towards integrated landscape and water management* (267–283). IWA Publishing.
- Ahern, J. (2013). Urban landscape sustainability and resilience: the promise and challenges of integrating ecology with urban planning and design. *Landscape Ecology* **28**(6): 1203–1212. Accessed April 25, 2015: <https://scholar-google-com.proxy.wexler.hunter.cuny.edu/scholar?oi=bibs&clus=ter=2360019842460228842&btnI=1&hl=en>
- Akbari, H., Pomerantz, M., and Taha, H. (2001). Cool surfaces and shade trees to reduce energy use and improve air quality in urban areas. *Solar Energy* **70**(3), 295–310.
- Alberti, M. (2008). *Advances in Urban Ecology: Integrating Humans and Ecological Processes in Urban Ecosystems*. Springer Science + Business Media.
- Alberti, M. (2015). Eco-evolutionary dynamics in an urbanizing planet. *TREE*, in press. Accessed January 8, 2016: <http://dx.doi.org/10.1016/j.tree.2014.11.007>
- Alberti, M., and Marzluff, J. M. (2004). Ecological resilience in urban ecosystems: Linking urban patterns to human and ecological functions. *Urban Ecosystems* **7**, 241–265.
- Alexandri, E., and Jones, P. (2008). Temperature decreases in an urban canyon due to green walls and green roofs in diverse climates. *Building and Environment*, **43**, 480–493. Accessed March 2, 2014: <http://www.sciencedirect.com/science/article/pii/S0360132306003957>
- Amin, A. (2007). Re-thinking the urban social. *City* **11**(1), 100–114.
- Andersson, E., Barthel, S., Borgström, S., Colding, J., Elmqvist, T., Folke, C., and Gren, Å. (2014). Reconnecting cities to the biosphere: Stewardship of green infrastructure and urban ecosystem services. *Ambio* **43**, 445–453. Accessed February 27, 2015: <http://dx.doi.org/10.1007/s13280-014-0506-y>.
- Andersson, E., McPhearson, T., Kremer, P., Gomez-Baggethun, E., Haase, D., Tuvaldal, M., and Wurster, D. (2015a). Scale and Context Dependence of Ecosystem Service Providing Units. *Ecosystem Services* (Special Issue). Accessed March 25, 2016: <http://dx.doi.org/10.1016/j.ecoser.2014.08.001>
- Andersson, E., Tengö, M., McPhearson, T., and Kremer, P. (2015b). Cultural ecosystem services as a platform for working towards urban sustainability. *Ecosystem Services* (Special Issue). Accessed February 6, 2016: <http://dx.doi.org/10.1016/j.ecoser.2014.08.002>
- Aronson, M. F. J., La Sort, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C. A. et al. (2014). A global analysis of the impacts of urbanization on bird and plant diversity reveals key anthropogenic drivers. *Proceedings of the Royal Society B* **281**(2013), 3330.
- Asian Cities Climate Change Resilience Network. (2015). Urban Climate Change Resilience in Action: Lessons from Projects in 10 ACCCRN Cities. ACCCRN & The Rockefeller Foundation. Accessed March 30, 2016: http://accrn.net/sites/default/files/publication/attach/ACCCRN_ProjectsInsightsPaper.pdf
- Asikainen, E., and Jokinen, A. (2009). Future natures in the making: Implementing biodiversity in sub-urban land use planning. *Planning Theory and Practice* **10**(3), 351–368.
- Bardsley, D. K., and Hugo, G. J. (2010). Migration and climate change: Examining thresholds of change to guide effective adaptation decision-making. *Population and Environment* **32**, 238–262. doi:10.1007/s11111-010-0126-9
- Barthel, S., and Isendahl, C. (2013). Urban gardens, agriculture, and water management: Sources of resilience for long-term food security in cities. *Ecological Economics* **86**(2), 224–234.

- Barthel, S., Parker, J., and Ernstson, H. (2015). Food and green space in cities: A resilience lens on gardens and urban environmental movements. *Urban Studies* **52**(7), 1321–1338.
- Batty, M. (2008). The size, scale, and shape of cities. *Science* **319**, 769–771. doi:10.1126/science.1151419
- Beatley, T. (2000). *Green Urbanism*. Island Press.
- Beatley, T. (2009). *Biophilic Cities*. Island Press.
- Berkes, F., and Folke, C. (1998). *Linking Social and Ecological Systems: Management Practices and Social Mechanisms for Building Resilience*. Cambridge University Press.
- Bettencourt, L., and West, G. (2010). A unified theory of urban living. *Nature*, **467**, 912–913. doi:10.1038/467912a.
- Bianchini, F., and Hewage, K. (2012). Probabilistic social cost-benefit analysis for green roofs: A lifecycle approach. *Building and Environment* **58**, 152–162. doi:10.1016/j.buildenv.2012.07.005
- Bolund, P., and Hunhammar, S. (1999). Ecosystem services in urban areas. *Ecological Economics* **29**, 293–301. doi:10.1016/S0921-8009(99)00013-0
- Boone, C. G., Buckley, G. L., Grove, J. M., and Sister, C. (2009). Parks and people: An environmental justice inquiry in Baltimore, Maryland. *Annals of the Association of American Geographers* **99**(July 2008), 767–787.
- Booth, D.B., Haugerud, R.A., and Troost, K.G. 2003. Geology, watershed, and Puget lowland rivers. In Montgomery, D.R., Bolton, S., Booth, D.B. and Wall, L. (Eds.) *Restoration of Puget Sound Rivers*. The University of Washington Press. Seattle, WA. pp. 14–45.
- Boosabong, P. (2014). Coping with flooding in Bangkok. *Urban Agriculture Magazine* **27**.
- Borgström, S. T., Elmquist, T., Angelstam, P., and Alfsen-Norodom, C. (2006). Scale mismatches in management of urban landscapes. *Ecology and Society* **11**(2), 16. doi:10.1097/MCC.0b013e32807f2aa5
- Braaker, S., Moretti, M., Boesch, R., Ghazoul, J., Obrist, M. K., and Bontadina, F. (2014). Assessing habitat connectivity for ground-dwelling animals in an urban environment. *Ecological Applications* **24**, 1583–1595.
- Breuste, J., Niemelä, J., and Snep, R. P. H. (2008). Applying landscape ecological principles in urban environments. *Landscape Ecology* **23**(10), 1139–1142. doi:10.1007/s10980-008-9273-0
- Cardinale, B. J., Duffy, J. E., Gonzalez, A., Hooper, D. U., Perrings, C., Venail, P., and Naeem, S. (2012). Biodiversity loss and its impact on humanity. *Nature* **486**(7401), 59–67. doi:10.1038/nature11148
- Carson, T. B., Marasco, D. E., Culligan, P. J., and McGillis, W. R. (2013). Hydrological performance of extensive green roofs in New York City: Observations and multi-year modelling of three full-scale systems. *Environmental Research Letters* **8**(02), 4036. doi:10.1088/1748-9326/8/2/024036
- CDKN. (2014). Inside stories on climate compatible development. Climate Development Knowledge Network (CDKN) Accessed December 21, 2015: http://cdkn.org/wp-content/uploads/2014/05/SriLanka_Inside_Story_final_web-res.pdf
- Cerveny, R. S., and Balling, R. C., Jr. (1998). Weekly cycles of air pollutants, precipitation and tropical cyclones in the coastal NW Atlantic region. *Nature* **394**, 561–563. doi:10.1038/29043
- Chan, K. M., Satterfield, T., and Goldstein, J. (2011). Rethinking ecosystem services to better address and navigate cultural values. *Ecological Economics* **74**, 8–18. doi:10.1016/j.ecolecon.2011.11.011
- Childers, D.L., Cadenasso, M.L., Grove, J.M., Marshall, V., MacGrath, B., and Pickett, S. 2015. An ecology for cities: A transformational nexus of design and ecology to advance climate change resilience and urban sustainability. *Sustainability* **7**:3774–3791.
- Childers, D. L., Pickett, S. T. A., Grove, J. M., Ogden, L., and Whitmer, A. (2014). Advancing urban sustainability theory and action: Challenges and opportunities. *Landscape & Urban Planning* **125**, 320–328.
- Cohen, N., and Wijsman, K. (2014). Urban agriculture as green infrastructure: The case of New York City. *Urban Agriculture Magazine* **27**.
- Colding, J. (2011). The role of ecosystem services in contemporary urban planning. In Niemela, J. (ed.), *Urban Ecology*. Oxford University Press, 228–237.
- Colding, J. (2013). Local assessment of Stockholm: Revisiting the Stockholm Urban Assessment. In Elmquist, T., Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P. J., McDonald, R. I., Parnell, S., Schewenius, M., et al. (eds.), *Urbanization, Biodiversity and Ecosystem Services – Challenges and Opportunities* (313–335). Springer Netherlands. doi:10.1007/978-94-007-7088-1
- Cole, L. W., and Foster, S. R. (2001). *From the Ground Up: Environmental Racism and the Rise of the Environmental Justice Movement* (244). NYU Press.
- Collins, J. P., Kinzig, A., Grimm, N. B., Fagan, W. F., Hope, D., Wu, J., and Borer, E. T. (2000). A new urban ecology. *American Scientist* **88**, 416–425.
- Condon, P. M., Cavens, D., and Miller, N. (2009). Urban planning tools for climate change mitigation. *Policy focus report/Code PF021*. Accessed May 28, 2015: www.lincolnst.edu
- Consolo, T., Faux, M., Becker, M., Hamstead, Z., Kremer, P., Malone, E., and Segal, P. (2013). Equitable re-development of land acquired by New York City and New York State in Sandy-affected areas. Planner Network. Accessed March 13, 2014: <http://596acres.org/en/news/2013/12/19/what-our-new-may-or-should-do-redevelopment-in-sandy-affected-communities/>
- Convention on Biological Diversity (CBD). (2009). Connecting biodiversity and climate change mitigation and adaptation: Report of the second ad hoc Technical Expert Group on Biodiversity and Climate Change. Technical series No. 41. Secretariat of the Convention on Biological Diversity, Montreal.
- Corburn, J. (2005). *Street Science*. MIT Press.
- Cumming, G. S., Cumming, D. H. M., and Redman, C. L. (2006). Scale mismatches in social-ecological systems: Causes, consequences, and solutions. *Ecology and Society* **11**(1), 20.
- Czemieli Berndtsson, J. (2010). Green roof performance towards management of runoff water quantity and quality: A review. *Ecological Engineering* **36**, 351–360. doi:10.1016/j.ecoleng.2009.12.014
- Dale, A. G., and Frank, S. D. (2014). Urban warming trumps natural enemy regulation of herbivorous pests. *Ecological Applications* **24**, 1596–1607.
- De la Salle, J., and Holland, M. (2010). *Agricultural Urbanism: Handbook for Building Sustainable Food Systems in 21st Century Cities*. Green Frigate Books.
- Demuzere, M., Orru, K., Heidrich, O., Olazabal, E., Geneletti, D., Orru, H., Bhave, A. G., and Faehnle, M. (2014). Mitigating and adapting to climate change: Multi-functional and multi-scale assessment of green urban infrastructure. *Journal of Environmental Management* **146**, 107–115.
- Depietri, Y., Renaud, F. G., and Kallis, G. (2012). Heat waves and floods in urban areas: A policy-oriented review of ecosystem services. *Sustainability Science* **7**, 95–107. doi:10.1007/s11625-011-0142-4
- De Zeeuw, H., van Veenhuizen R., and Dubbeling M. (2011). The role of urban agriculture in building resilient cities in developing countries. *Journal of Agricultural Science* **149**(S1), 153–163.
- Di Leo, N., Escobedo, F. J., and Dubbeling, M. (2015). The role of urban green infrastructure in mitigating land surface temperature in Bobo-Dioulasso, Burkina Faso. *Environment Development and Sustainability*. doi:10.1007/s10668-015-9653-y.
- Dubbeling, M. (2014). Integrating urban agriculture and forestry into climate change action plans: Lessons from Sri Lanka. Accessed November 26, 2015: <http://www.ruaf.org/publications/integrating-urban-agriculture-and-forestry-climate-change-action-plans-lessons-sri-0>
- Dubbeling, M. (2015). Integrating urban agriculture and forestry into climate change action plans: Lessons from Western Province, Sri Lanka and Rosario, Argentina. RUAF Foundation.
- Dunnett, N., and Kingsbury, N. (2008). *Planting Green Roofs and Living Walls*. Timber Press.
- Ebre Observatory. (2014). Accessed March 11, 2015: <http://www.obsebre.es/en/>
- Ecologic Institute. (2011). Analysis of socio-economic driving forces in built-up area expansion in Xiamen, China. In *Issues in Ecological Research and Application: 2011 Edition*. Scholarly Editions.
- Eisen, L., Monaghan, A. J., Lozano-Fuentes, S., Steinhoff, D. F., Hayden, M. H., and Bieringer, P. E. (2014). The impact of temperature on the

- biomics of *Aedes (Stegomyia) aegypti*, with special reference to the cool geographic range margins. *Journal of Medical Entomology* **51**(3), 496–516. doi:<http://dx.doi.org/10.1603/ME13214>
- Elmqvist, T., Redman, C. E., Barthel, S., and Costanza, R. (2013). History of urbanization and the missing ecology. In Elmqvist, T., Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P. J., McDonald, R. I., and Wilkinson, C. (eds.), *Urbanization, Biodiversity and Ecosystem Services: Challenges and Opportunities* (13–30). Springer. doi:10.1007/978-94-007-7088-1_2
- Ernstson, H. (2013). The social production of ecosystem services: A framework for studying environmental justice and ecological complexity in urbanized landscapes. *Landscape and Urban Planning* **109**(1), 7–17. doi:10.1016/j.landurbplan.2012.10.005
- Ernstson, H., Barthel, S., Andersson, E., and Borgström, S. T. (2010). Scale-crossing brokers and network governance of urban ecosystem services: The case of Stockholm. *Ecology and Society* **15**(4), 28. Accessed October 24, 2015: <http://www.ecologyandsociety.org/vol15/iss4/art28/main.html>
- Escobedo, F. J., Clerici, N., Staudhammer, C. L., and Corzo, G. T. (2015). Socio-ecological dynamics and inequality in Bogotá, Colombia's public urban forests and their ecosystem services. *Urban Forestry & Urban Greening* **14**(4), 1040–1053.
- European Environmental Agency. (2015). Exploring nature-based solutions – The role of green infrastructure in mitigating the impacts of weather- and climate change-related natural hazards. EEA Technical Report No. 12.
- FAO. (2014). Growing greener cities in Latin America and the Caribbean. A FAO report on urban and peri-urban agriculture in the region. Rome, Food and Agriculture Organization of the United Nations.
- Farr, D. (2008). *Sustainable Urbanism: Urban Design With Nature*. John Wiley & Sons.
- Farrell, C., Szota, C., and Arndt, S. K. (2015). Urban plantings: “Living laboratories” for climate change response. *Trends in Plant Science* **20**(10), 597–599.
- Fitzpatrick, M. C., Gove, A. D., Sanders, N. J., and Dunn, R. R. (2008). Climate change, plant migration, and range collapse in a global biodiversity hotspot: The *Banksia* (Proteaceae) of Western Australia. *Global Change Biology* **14**, 1337–1352.
- Florgård, C. (2007). Treatment measures for original natural vegetation preserved in the urban green infrastructure at Jarvafaltet, Stockholm. In Stewart, M., Ignatieva, M., Bowring, J., Egoz, S., and Melnichuk, I. (eds.), *Globalisation of Landscape Architecture: Issues for Education and Practice* (100–102). St. Petersburg's State Polytechnic University Publishing House.
- Folke, C., Carpenter, S., Walker, B., Scheffer, M., Elmqvist, T., Gunderson, L., and Holling, C. S. (2004). Regime shifts, resilience, and biodiversity in ecosystem management. *Annual Review of Ecology, Evolution, and Systematics* **35**, 557–581.
- Folke, C. (2010). How resilient are ecosystems to global environmental change? *Sustainability Science* **5**, 151–154. doi:10.1007/s11625-010-0109-x
- Fook, L. L., and Gang, C. (eds.). (2010). *Towards a Liveable and Sustainable Urban Environment: Eco-Cities in East Asia*. World Scientific Publishing.
- Foster, J., A. Lowe, and S. Winkelman. (2011). *The Value of Green Infrastructure for Urban Climate Adaptation*. Washington, D.C.: Center for Clean Air Policy. Accessed May 5, 2015: http://mi.mi.gov/documents/dnr/TheValue_347538_7.pdf.
- Fowler, D. (2002). Pollutants deposition and uptake by vegetation. In Bell, J. N. B., and Treshow, M. (eds.), *Air Pollution and Plant Life*, second edition (43). Wiley.
- Frantzeskaki, N., and Tilie, N. (2014). The dynamics of urban ecosystem governance in Rotterdam, The Netherlands. *Ambio* **43**(4), 542–555. Accessed August 12, 2015: <http://doi.org/10.1007/s13280-014-0512-0>
- Frantzeskaki, N., Kabisch, N., and McPhearson, T. (2016). Advancing urban environmental governance: Understanding theories, practices and processes shaping urban sustainability and resilience. *Environmental Science and Policy* (Special Issue) **62**, 1–6. Accessed May 18, 2017: <http://dx.doi.org/10.1016/j.envsci.2016.05.008>
- Frischenbruder, M., and Pellegrino, P. (2006). Using greenways to reclaim nature in Brazilian cities. *Landscape and Urban Planning* **76**, 67–78.
- Frumkin, H., Hess, J., Luber, G., Malilay, J., and McGeehin, M. (2008). Climate change: The public health response. *American Journal of Public Health* **98**(3), 435–445. doi:10.2105/AJPH.2007.119362
- Georgescu, M., Morefield, P. E., Bierwagen, B. G., and Weaver, C. P. (2014). Urban adaptation can roll back warming of emerging megapolitan regions. *Proceedings of the National Academy of Science U.S.A.* **111**, 2909–2914. doi:10.1073/pnas.1322280111.
- Gill, S. E., Handley, J. F., Ennos, A. R., and Pauleit, S. (2007). Adapting cities for climate change: The role of the green infrastructure. *Built Environment* **33**(1), 115–133.
- Gill, S. E., Rahman, M. A., Handley, J. F., and Ennos, A. R. (2013). Modelling water stress to urban amenity grass in Manchester UK under climate change and its potential impacts in reducing urban cooling. *Urban Forestry & Urban Greening* **12**, 350–358. Accessed June 4, 2014: <http://dx.doi.org/10.1016/j.ufug.2013.03.005>
- Gilman, Sarah E., Urban, M. C., Tewksbury, J., Gilchrist, G. W., and Holt, R. D. (2010). A framework for community interactions under climate change. *Trends in Ecology and Evolution* **25**, 325–331. Accessed April 25, 2014: doi:10.1016/j.tree.2010.03.002
- Gillner, S., Bräuning, A., and Roloff, A. (2014). Dendrochronological analysis of urban trees: climatic response and impact of drought on frequently used tree species. *Trees* **28**, 1079. doi:10.1007/s00468-014-1019-9
- Giroi, P., Ehrhart, C., and Oglethorpe, J. (2009). *Integrating Community and Ecosystem-Based Approaches in Climate Change Adaptation Responses*. Ecosystem and Livelihoods Adaptation Network (ELAN).
- Glick, P., Stein, B. A., and Edelson, N. A. (eds.). (2011). *Scanning the Conservation Horizon: A Guide to Climate Change Vulnerability Assessment*. National Wildlife Federation.
- Gómez-Baggethun, E., Gren, Å., Barton, N. D., Langemeyer, J., McPhearson, T., O'Farrell, P., and Kremer, P. (2013). Urban ecosystem services. In Elmqvist, T., Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P. J., McDonald, R. I., and Wilkinson, C. (eds.), *Urbanization, Biodiversity and Ecosystem Services: Challenges and Opportunities* (175–251). Springer Netherlands. doi:10.1007/978-94-007-7088-1
- Gonzalez, P., R.P. Nielson, R.P., Lenihan, J.M., and Drapek, R.J. 2010. Global patterns in the vulnerability of the ecosystems to vegetation shifts due to climate change. *Global Ecology and Biogeography* **19**:755–768.
- Gotelli, N. J., and Chao, A. (2013). Measuring and estimating species richness, species diversity, and biotic similarity from sampling data. In Levin, S. A. (ed.), *Encyclopedia of Biodiversity*, second edition, volume **5**(195–211). Academic Press. Accessed November 27, 2015: <http://www.elsevier.com/locate/permissions>
- Grimm, N. B., Chapin, F. S., Bierwagen, B., Gonzalez, P. Groffman, P. M., Luo, Y., Melton, F., Nadelhoffer, K., Pairis, A., Raymond, P. A., Schimel, J., and Williamson, C. E. (2013). The impacts of climate change on ecosystem structure and function. *Frontiers in Ecology and the Environment* **11**, 474–482.
- Grimm, N. B., Foster, D., Groffman, P., Grove, J. M., Hopkinson, C. S., Nadelhoffer, K. J., Peters, D. P. (2008). The changing landscape: Ecosystem responses to urbanization and pollution across climatic and societal gradients. *Frontiers in Ecology and the Environment* **6**(5), 264–272. doi:10.1890/070147
- Grimm, N. B., Grove, J. M., Pickett, S. T. A., and Redman, C. A. (2000). Integrated approaches to long-term studies of urban ecological systems. *BioScience* **50**(7), 571. doi:10.1641/0006-3568(2000)050[0571:IALTO]2.0.CO;2
- Güneralp, B., McDonald, R. I., Fragkias, M., Goodness, J., Marcotullio, P. J., and Seto, K. C. (2013). Urbanization, biodiversity and ecosystem services: Challenges and opportunities. In Elmqvist, Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P. J., McDonald, R. I., and Wilkinson, C. (eds.), *Urbanization, Biodiversity and Ecosystem Services: Challenges and Opportunities* (437–452). Springer Netherlands. doi:10.1007/978-94-007-7088-1
- Güneralp, B., and Seto, K. C. (2013). Futures of global urban expansion: Uncertainties and implications for biodiversity conservation. *Environmental Research Letters* **8**(1), 014025. doi:10.1088/1748-9326/8/1/014025

- Haase, D., Haase, A., Kabisch, N., Kabisch, S., and Rink, D. (2012). Actors and factors in land-use simulation: *The challenge of urban shrinkage*. *Environmental Modelling and Software* **35**, 92–103. doi:10.1016/j.envsoft.2012.02.012
- Haase, D., Larondelle, N., Andersson, E., Artmann, M., Borgström, S., Breuste, J., Gomez-Baggethun, E., Gren, Å., Hamstead, Z., Hansen, R., Kabisch, N., Kremer, P., Langemeyer, J., Rall, E. L., McPhearson, T., Pauleit, S., Qureshi, S., Schwarz, N., Voigt, A., Wurster, D., and Elmqvist, T. (2014). A quantitative review of urban ecosystem services assessments: Concepts, models, and implementation. *AMBIO* (Special Issue) **43**, 413–433. doi:10.1007/s13280-014-0504-0
- Hallegatte, S., Ranger, N., Mestre, O., Dumas, P., Corfee-Morlot, J., Herweijer, C., and Wood, R. M. (2010). Assessing climate change impacts, sea level rise and storm surge risk in port cities: A case study on Copenhagen. *Climatic Change* **104**(1), 113–137. doi:10.1007/s10584-010-9978-3
- Hatfield, J. L., Boote, K. J., Kimball, B. A., Ziska, L. H., Izaurralde, R. C., Ort, D., Thomson, A. M., and Wolfe, D. W. (2011). Climate impacts on agriculture: Implications for crop production. *Agronomy Journal* **103**(2011), 351–370.
- Hamstead, Z. A., Larondelle, N., Kremer, P., McPhearson, T., and Haase, D. (2015). Classification of the heterogeneous structure of urban landscapes (STURLA) as an indicator of landscape function applied to surface temperature in New York City. *Ecological Indicators* (Special Issue). Accessed April 12, 2016: <http://dx.doi.org/10.1016/j.ecolind.2015.10.014>
- Hansen, R., Frantzeskaki, N., McPhearson, T., Rall, E., Kabisch, N., Kaczorowska, A., Kain, J. -H., Artmann, M., and Pauleit, S. (2015). The uptake of the ecosystem services concept in planning discourses of European and American cities. *Ecosystem Services* (Special Issue). doi:10.1016/j.ecoser.2014.11.013
- Heller, N. E., and Zavaleta, E. S. (2009). Biodiversity management in the face of climate change: A review of 22 years of recommendations. *Biological Conservation* **142**, 14–32.
- Heutte, R., Bella, E., Snyder, J., and Shephard, M. 2003. Invasive plants and exotic weeds of Southeast Alaska. *USDA Forest Service*. Anchorage.
- Hunt, A., and Watkiss, P. (2011). Climate change impacts and adaptation in cities: A review of the literature. *Climatic Change* **104**, 13–49.
- Huq, N., Renaud, F., and Sebesvari, Z. (2013). *Ecosystem Based Adaptation (EBA) to Climate Change – Integrating Actions to Sustainable Adaptation*. United Nations University, Institute for Environment and Human Security (UNU-EHS), UN Campus, Bonn, Germany.
- ICLEI Africa Secretariat 2013 *Annual Report*. (2013). ICLEI Local Governments for Sustainability.
- ICIMOD. (2009). *Local Responses to Too Much and Too Little Water in the Greater Himalayan Region*. ICIMOD. Accessed September 15, 2014: <http://lib.icimod.org/record/8021>
- Ignatieva, M., Stewart, G.H., and Meurk, C. (2011). Planning and design of ecological networks in urban areas. *Landscape Ecol Eng.* **7**:17–25.
- Intergovernmental Panel on Climate Change (IPCC). (2001). *Climate Change 2001: The Scientific Basis. A Report of Working Group I of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Intergovernmental Panel on Climate Change (IPCC). (2007a). *Climate Change 2007: Synthesis Report*. Cambridge University Press.
- Intergovernmental Panel on Climate Change (IPCC). (2013). Summary for policymakers. In Stocker, T. F., Qin, D., Plattner, G. K., Tignor, M., Allen, S. K., Boschung, J., and Midgley, P. M. (eds.), *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Intergovernmental Panel on Climate Change (IPCC). (2014). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. IPCC Working Group II Contribution to AR5*. Accessed August 13, 2015: <http://ipcc-wg2.gov/AR5/>
- Institute for Social and Environmental Transition (ISET). (2010). *The Shared Learning Dialogue: Building Stakeholder Capacity and Engagement for Resilience Action*. Resilience in Concept and Practice Series Working Paper No. 1. ISET.
- IUCN. (2009). *Ecosystem-based Adaptation (EBA): Policy Briefing. Fifth session of the UNFCCC Ad Hoc Working Group on Long-Term Cooperative Action under the Convention (AWG-LCA)*, March 29–April 8, 2009, Bonn.
- Jansma, J. E., Sukkel, W., Stilma, E. S. C., vanOost, A. C. J., and Visser, A. J. (2012). The impact of local food production on food miles, fossil energy use and greenhouse gas (GHG) emission: The case of the Dutch city of Almere. In Viljoen, A., and Wiskerke, J. S. C. (eds.), *Sustainable Food Planning: Evolving Theory and Practice* (307–321). Wageningen Academic Publishers.
- Karki, M.B., Shrestha, A.B., and Winiger, M. (2011). Enhancing Knowledge Management and Adaptation Capacity for Integrated Management of Water Resources in the Indus River Basin. *Mountain Research and Development*. **31**(3):242–251.
- Kaushal, S. S., Groffman, P. M., Band, L. E., Shields, C. A., Morgan, R. P., Palmer, M. A., Belt, K. T., Swan, C. M., Findlay, S. E. G., and Fisher, G. T. (2008). Interaction between urbanization and climate variability amplifies watershed nitrate export in Maryland. *Environmental Science and Technology* **42**, 5872–5878.
- Kaye, J. P., Burke, I. C., Mosier, A. R., and Guerschman, J. P. (2004). Methane and nitrous oxide fluxes from urban soils to the atmosphere. *Ecological Applications* **14**, 975–981.
- Keim, M. E. (2008). Building human resilience: The role of public health preparedness and response as an adaptation to climate change. *American Journal of Preventive Medicine* **35**(5), 508–16. doi:10.1016/j.amepre.2008.08.022
- Kendal, D., Williams, N. S. G., and Williams, K. J. H. (2012). A cultivated environment: Exploring the global distribution of plants in gardens, parks and streetscapes. *Urban Ecosystems* **15**, 637–652. doi:10.1007/s11252-011-0215-2
- Kendle T. and Forbes S. 1997. Urban nature conservation. *E. & F.N. Spon*. London.
- Kenward, A., et al. (2014). *Summer in the City: Hot and Getting Hotter*. Climate Central. Accessed November 27, 2015: <http://www.climatecentral.org/>
- Kingsolver, J. G., Diamond, S. E., and Buckley, L. B. (2013). Heat stress and the fitness consequences of climate change for terrestrial ectotherms. *Functional Ecology*. **27**(6):1415–1423. <https://scholar.google.com/scholar?oi=bibs&cluster=10819040795748572617&btnI=1&hl=en>
- Klausmeyer, K. R., Shaw, R., MacKenzie, J. B., and Cameron, D. R. (2011). Landscape-scale indicators of biodiversity's vulnerability to climate change. *Ecosphere* **2**(8), art88. Accessed May 5, 2014: [http://dx.doi.org/10.1890/ES11-0\(0044\).1](http://dx.doi.org/10.1890/ES11-0(0044).1)
- Kopperoinen, L., Itkonen, P., and Niemelä, J. (2014). Using expert knowledge in combining green infrastructure and ecosystem services in land use planning: An insight into a new place-based methodology. *Landscape Ecology* **29**, 1361–1375. doi:10.1007/s10980-014-0014-2
- Kousky, C., Olmstead, S., Walls, M., and Macauley, M. (2013). Strategically placing green infrastructure: Cost-effective land conservation in the floodplain. *Environmental Science & Technology* **47**(8), 3563–3570.
- Kowarik, I. (2005) Wild Urban Woodlands: Towards a Conceptual Framework. *Wild Urban Woodlands*. 1–32.
- Kremer, P., Hamstead, Z., Haase, D., McPhearson, T., Frantzeskaki, N., Andersson, E., Kabisch, N., Larondelle, N., Lorange Rall, E., Avlonitis, G., Bertram, C., Baró, F., Gómez-Baggethun, E., Gren, A., Hansen, R., Kaczorowska, A., Kain, J. -H., Kronenberg, J., Langemeyer, J., Muehlmann, P., Pauleit, S., Rehdanz, K., Schewenius, M., van Ham, C., Voigt, A., Wurster, D., and Elmqvist, T. (2016a). Key insights for the future of urban ecosystem services research. *Ecology and Society* **21**(2), 29. Accessed January 12, 2017: <http://dx.doi.org/10.5751/ES-08445-210229>
- Kremer, P., Hamstead, Z. A., and McPhearson, T. (2016b). The value of urban ecosystem services: A spatially explicit multicriteria analysis of landscape scale valuation scenarios in NYC. *Environmental Science and Policy* (Special Issue). Accessed February 15, 2017: <http://dx.doi.org/10.1016/j.envsci.2016.04.012>
- Kronenberg, J. (2015). Why not to green a city? Institutional barriers to preserving urban ecosystem services. *Ecosystem Services*. **12**:218–227. <https://scholar-google-com.proxy.wexler.hunter.cuny.edu/scholar?oi=bibs&cluster=3678534514957264970&btnI=1&hl=en>

- Larondelle, N., Hamstead, Z. A., Kremer, P., Haase, D., and McPhearson, T. (2014). Applying a novel urban structure classification to compare the relationships of urban structure and surface temperature in Berlin and New York City. *Applied Geography* 53, 427–437. Accessed February 7, 2015: <http://dx.doi.org/10.1016/j.apgeog.2014.07.004>
- Laros, M., Birch, S., and Clover, J. (2013). *Ecosystem-Based Approaches to Building Resilience in Urban Areas: Towards a Framework for Decision-Making Criteria*. Background paper. ICLEI-Africa.
- La Rosa, D., and Privitera, R. (2013). Characterization of non-urbanized areas for land-use planning of agricultural and green infrastructure in urban contexts. *Landscape and Urban Planning* 109(1), 94–106. doi:10.1016/j.landurbplan.2012.05.012
- Leichenko, R. M., and O'Brien, K. L. (2008). *Environmental Change and Globalization: Double Exposures*. Oxford University Press.
- Leipprand, A., and Gerten, D. (2006). Global effects of doubled atmospheric CO₂ content on evapotranspiration, soil moisture and runoff under potential natural vegetation. *Hydrological Sciences Journal* 51, 171–185.
- Loarie, S. R., Carter, B. E., Hayhoe, K., McMahon, S., Moe, R., Knight, C. A., and Ackerly, D. D. (2008). Climate change and the future of California's endemic flora. *PLoS ONE* 3, e2502.
- Locatelli, B., Brockhaus, M., Buck, A., and Thompson, I. (2010). Forests and adaptation to climate change: Challenges and opportunities. In Mery, G., et al. (eds.), *Forest and Society: Responding to Global Drivers of Change*. IUFRO World Series.
- Loukaitou-sideris, A., and Stieglitz, O. (2002). Children in Los Angeles parks. *Town Planning Review* 73, 4(April), 467–488.
- Lye, L.H. (2010) The Judiciary and Environmental Governance in Singapore. *Journal of Court Innovation*. 3:133.
- Lwasa, S., and Dubbeling, M. (2015). Urban agriculture and climate change. In de Zeeuw, H., and Dreschel, P. (eds.), *Cities and Agriculture: Developing Resilient Urban Food Systems*. Earthscan-Routledge.
- Lwasa, S., Mugaga, F., Wahab, B., Simon, D., Connors, J., and Griffith, C. (2013). Urban and peri-urban agriculture and forestry: Transcending poverty alleviation to climate change mitigation and adaptation. *Urban Climate*. doi:10.1016/j.uclim.2013.10.007
- Maimaitiyiming, M., et al. (2014). Effects of green space spatial pattern on land surface temperature: Implications for sustainable urban planning and climate change adaptation. *Journal of Photogrammetry and Remote Sensing* 89, 59–66. Accessed December 27, 2015: <http://dx.doi.org/>
- Marcotullio, P. J., and Solecki, W. D. (2013). Sustainability and cities – Meeting the grand challenge for the twenty-first century. In Sygna, L., O'Brien, K., and Wolf, J. (eds.), *A Changing Environment for Human Security: Transformative Approaches to Research, Policy and Action* (496). Routledge.
- Marzluff, J. M. (2001). Worldwide urbanization and its affects on birds. In Marzluff, J. M., Bowman, R., and Donnelly, R. (eds.), *Avian Conservation and Ecology in an Urbanizing World* (19–47). Kluwer Academic Publishers.
- McDonald, R., and Marcotullio, P. (2011). Global effects of urbanization on ecosystem services. In Niemelä, J. (ed.), *Urban Ecology* (193–205). Oxford University Press.
- McDonald, R. I., Marcotullio, P. J., and Güneralp, B. (2013). Urbanization and global trends in biodiversity and ecosystem services. In Elmquist, T., Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P. J., McDonald, R. I., and Wilkinson, C. (eds.), *Urbanization, Biodiversity and Ecosystem Services: Challenges and Opportunities* (31–52). Springer Netherlands. doi:10.1007/978-94-007-7088-1_3
- McDonnell, M. J. (2011). The history of urban ecology. In Niemelä J., Breuste, J. H., Guntenspergen, G., McIntyre, N. E., Elmquist, T., and James, P. (eds.), *Urban Ecology: Patterns, Processes, and Applications*. Oxford University Press.
- McDonnell, M. J., and Hahs, A. (2013). The future of urban biodiversity research: Moving beyond the “low-hanging fruit.” *Urban Ecosystems* 16, 397–409. doi:10.1007/s11252-013-0315-2
- McGranahan, G., Marcotullio, P. J., Bai, X., Balk, D., Braga, T., Douglas, I., and Zlotnik, H. (2005). Urban systems. In Hassan, R., Scholes, R., and Ash, N. (eds.), *Current State and Trends: Findings of the Condition and Trends Working Group. Ecosystems and Human Well-being* (795–825). Island Press.
- McKinney, M. L. (2002). Urbanization, Biodiversity and Conservation. *BioScience* 52(10), 883–890.
- McLennan, J. F. (2004). *The Philosophy of Sustainable Design: The Future of Architecture*. Ecotone Publishing. Accessed February 22, 2014: http://books.google.com/books?id=Qjadh_0leMC&pgis=1
- McPhearson, T. (2011). Toward a sustainable New York City: Greening through urban forest restoration. In Slavin, M. (ed.), *The Triple Bottom Line: Sustainability Principles, Practice, and Perspective in America's Cities* (181–204). Island Press.
- McPhearson, T., Auch, R., and Alberti, M. (2013c). Urbanization trends, biodiversity patterns, and ecosystem services in North America. In Elmquist, T., Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P. J., McDonald, R. I., Parnell, S., Schewenius, M., Sendstad, M., Soto, K. C., and Wilkinson, C. (eds.), *Cities and Biodiversity Outlook: Urbanization, Biodiversity and Ecosystem Services: Challenges and Opportunities* (279–286). Springer Netherlands. doi:10.1007/978-94-007-7088-1_14
- McPhearson, T., Haase, D., Kabisch, N., Gren, Å. (2016). Advancing understanding of the complex nature of urban systems. *Ecological Indicators* 70, 566–573. Accessed April 18, 2017: <http://dx.doi.org/10.1016/j.ecolind.2016.03.054>
- McPhearson, T., Hamstead, Z., and Kremer, P. (2014). Urban ecosystem services for resilience planning and management in New York City. *AMBIO* 43, 502–515. doi:10.1007/s13280-014-0509-8
- McPhearson, T., Kremer, P., and Hamstead, Z. (2013). Mapping ecosystem services in New York City: Applying a social-ecological approach in urban vacant land. *Ecosystem Services* (2013), 11–26. Accessed October 29, 2015: <http://dx.doi.org/10.1016/j.ecoser.2013.06.005>
- McPhearson, T., Andersson, E., Elmquist, T., and Frantzeskaki, N. (2015). Resilience of and through urban ecosystem services. *Ecosystem Services*. 12:152–156. <http://www.sciencedirect.com/science/article/pii/S2212041614000837>
- McPhearson, T., Pickett, S. T. A., Grimm, N., Alberti, M., Elmquist, T., Niemelä, J., Weber, C., Haase, D., Breuste, J., and Qureshi, S. (2016). Advancing urban ecology toward a science of cities. *BioScience* 66(3), 198–212. doi:10.1093/biosci/biw002
- McPherson, E. G., Nowak, D., Heisler, G., Grimmond, S., Souch, C., Grant, R., and Rowntree, R. (1997). Quantifying urban forest structure, function, and value: The Chicago Urban Forest Climate Project. *Urban Ecosystems* 1(1), 49–61. doi:10.1023/A:1014350822458
- Meteo Estartit. (2015). Accessed January 5, 2016: <http://www.historique-meteo.net/europe/espagne/estartit/2015/09/>
- Meyer, H., Nillesen, A. L., and Zonneveld, W. (2012). Rotterdam: A city and a mainport on the edge of a delta. *European Planning Studies* 20, 71–94. doi:10.1080/09654313.2011.638498.
- Millennium Ecosystem Assessment. (2005). *Ecosystems and Human Well-Being: Millennium Ecosystem Assessment Scenarios for the Future of Ecosystem Services*. Island Press.
- Mincy, S. K., Hutten, M., Fischer, B. C., Evans, T. P., Stewart, S. I., and Vogt, J. M. (2013). Structuring institutional analysis for urban ecosystems: A key to sustainable urban forest management. *Urban Ecosystems* 16(3), 553–571. doi:10.1007/s11252-013-0286-3
- Ministry of Agriculture, Agrarian Development, Minor Irrigation, Industries, Environment, Arts and Cultural Affairs, Western Province. (2014). Climate change adaptation action plan (2015–2018). Western Province of Sri Lanka. Accessed September 18, 2015: http://www.ruaf.org/sites/default/files/Western%20Province%20Climate%20Change%20Action%20Plan_2015-2018.pdf
- Miyake, K. K., Maroko, A. R., Grady, K. L., Maantay, J. A., and Arno, P. S. (2010). Not just a walk in the park: Methodological improvements for determining environmental justice implications of park access in New York City for the promotion of physical activity. *Cities and the Environment* 3(1), 1–17. Accessed August 14, 2014: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3160641&tool=pmcentrez&rendertype=abstract>

- Mosely, S. (2014). Coastal cities and environmental change. *Environment and History* **20**, 517–533. doi:10.3197/096734014X14091313617280
- Mostafavi, M., and Doherty, G. (2010). *Ecological Urbanism*. Lars Muller Publishers.
- Müller, N. (2010). Most frequently occurring vascular plants and the role of non-native species in urban areas – A comparison of selected cities of the old and new worlds. In N. Müller, P. Werner, and J. G. Kelcey (Eds.), *Urban biodiversity and design* (pp. 227–242). Hoboken: Wiley-Blackwell.
- Munang, R., Thiaw, I., Alverson, K., Mumba, M., Liu, J. and Rivington, M. (2013). Climate change and Ecosystem-based Adaptation: a new pragmatic approach to buffering climate change impacts. *Current Opinion Environmental Sustainability*. 5:1–5.
- Munroe, R., Roe, D., Doswald, N., Spencer, T., Möller, I., Vira, B., Reid, H., Kontoleon, A., Giuliani, A., Castelli I., and Stephens, J. (2010). *Review of the Evidence Base for Ecosystem-based Approaches for Adaptation to Climate Change*. International Institute for Environment and Development (IIED).
- National Environmental Justice Advisory Council. (2002). Fish consumption and environmental justice. Accessed January 4, 2015: http://www.epa.gov/compliance/ej/resources/publications/nejac/fish-consump-report_1102.pdf
- New York State (NYS). (2013). New York State Community Development Block Grant Disaster Recovery (CDBG-DR) Program. Accessed April 12, 2016: <https://stormrecovery.ny.gov/housing/buyout-acquisition-programs>
- Niemelä, J., Breuste, J. H., Guntenspergen, G., McIntyre, N. E., Elmqvist, T., and James, P. (2011). *Urban Ecology: Patterns, Processes, and Applications*. Oxford University Press.
- Nobis, M. P., Jaeger, J. A. G., and Zimmerman, N. E. (2009). Neophyte species richness at the landscape scale under urban sprawl and climate warming. *Diversity and Distributions* **15**, 928–939. doi:10.1111/j.1472-4642.2009.00610.x
- Nowak, D. J., Hirabayashi, S., Bodine, A., and Hoehn, R. (2013). Modeled PM_{2.5} removal by trees in ten U.S. cities and associated health effects. *Environmental Pollution* **178**, 395–402. doi:10.1016/j.envpol.2013.03.050
- Novotny, V., Ahern, J., and Brown, P. (2010). *Water Centric Sustainable Communities: Planning, Retrofitting, and Building the Next Urban Environment*. Wiley.
- Observatori de l'Ebre. (2014). Accessed December 7, 2015: <http://www.obsebre.es/en/OECD/CDRF>.
- OECD/CDRF. (2009). *Trends in Urbanisation and Urban Policies in OECD Countries: What Lessons for China?* OECD/CDRF.
- Onishi, A., Cao, X., Ito, T., Shi, F., and Imura, H. (2010). Evaluating the potential for urban heat-island mitigation by greening parking lots. *Urban Forestry and Urban Greening* **9**(4), 323–332. doi:10.1016/j.ufug.2010.06.002
- Ottelé M., Van Bohemen H., Fraaij A. L. A. (2010). Quantifying the deposition of particulate matter on climber vegetation on living walls. *Ecological Engineering* **36**, 154–162.
- Pandey, R., and Bardsley, D. K. (2013). Human ecological implications of climate change in the Himalaya: Pilot studies of adaptation in agro-ecosystems within two villages from Middle Hills and Tarai, Nepal. In *Impacts World 2013, International Conference on Climate Change Effects*, Potsdam, May 27–30.
- Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology Evolution and Systematics* **37**, 637–669.
- Parmesan, C., and Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature* **421**(6918), 37–42.
- Pascual, U., Muradian, R., Rodríguez, L. C., and Duraiappah, A. (2010). Exploring the links between equity and efficiency in payments for environmental services: A conceptual approach. *Ecological Economics* **69**(6), 1237–1244. doi:10.1016/j.ecolecon.2009.11.004
- Pataki, D. E., Boone, C. G., Hogue, T. S., Jenerette, G. D., McFadden, J. P., and Pincetl, S. (2011). Ecohydrology bearings—invited commentary: Socio-ecohydrology and the urban water challenge. *Ecohydrology* **4**, 341–347. doi: 10.1002/eco.209
- Paul, M. J., and Meyer, J. L. (2001). Streams in the urban landscape. *Annual Review of Ecological Systems* **32**, 333.
- Pauleit, S., Liu, L., Ahern, J., and Kazmierczak, A. (2011). Multifunctional green infrastructure planning to promote ecological services in the city. In Niemelä, J. (ed.), *Urban Ecology* (272–286). Oxford University Press.
- Pearson, R. G., and Dawson, T. P. (2003). Predicting the impacts of climate change on the distribution of species: Are bioclimate envelope models useful? *Global Ecology & Biogeography* **12**, 361–371.
- Petkova, E. P., Bader, D. A., Anderson, G. B., Horton, R. M., Knowlton, K., and Kinney, P. L. (2014). Heat-related mortality in a warming climate: Projections for 12 U.S. cities. *International Journal of Environmental Research and Public Health* **11**(11), 11371–11383.
- Petralli, M., Prokopp, A., Morabito, M., Bartolini, G., Torrigiani, T., and Orlandini, S. (2006). Ruolo delle aree verdi nella mitigazione dell'isola di calore urbana: Uno studio nella città di Firenze. *Rivista Italiana Agrometeorologia* **1**, 51–58.
- Pham, T. T. H., Apparicio, P., Séguin, A. M., Landry, S., and Gagnon, M. (2012). Spatial distribution of vegetation in Montreal: An uneven distribution or environmental inequity? *Landscape and Urban Planning* **107**(3), 214–224. Accessed October 15, 2014: <http://dx.doi.org/10.1016/j.landurbplan.2012.06.002>
- Pickett, S., Burch W. R. Jr., Dalton, S. D. and Foresman, R. W. 1997. Integrated urban ecosystem research. *Urban Ecosyst.* **1**:183–84.
- Pickett, S., Cadenasso, M., Grove, J., Groffman, P., Band, L., Boone, C., Burch, W., Grimmond, S., Hom, J., Jenkins, J., Law, N., Nilon, C., Pouyat, R., Szlavecz, K., Warren, P., and Wilson, M. (2008). Beyond urban legends: An emerging framework of urban ecology, as illustrated by the Baltimore ecosystem study. *BioScience* **58**(2), 139–150.
- Pickett, S. T. A., Cadenasso, M. L., Grove, J. M., Nilon, C. H., Pouyat, R. V., Zipperer, W. C., and Costanza, R. (2001). Urban ecological systems: Linking terrestrial ecological, physical, and socioeconomic components of metropolitan areas. *Annual Review of Ecology and Systematics*, **32**, 127–57.
- Pielke, R. A., Marland, G., Betts, R. A., Chase, T. N., Eastman, J. L., Niles, J. O., and Running, S. W. (2002). The influence of land-use change and landscape dynamics on the climate system: Relevance to climate-change policy beyond the radiative effect of greenhouse gases. *Philosophical Transactions. Series A, Mathematical, Physical, and Engineering Sciences* **360**(1797), 1705–1719. doi:10.1098/rsta.2002.1027
- Porter, E. M., et al. (2013). Interactive effects of anthropogenic nitrogen enrichment and climate change on terrestrial and aquatic biodiversity. *Biogeochemistry* **114**(1–3), 93–120.
- Porter, J. R., Dyball, R., Dumaresq, D., Deutsch, L., and Matsuda, H. (2014). Feeding capitals: Urban food security and self-provisioning in Canberra, Copenhagen and Tokyo. *Global Food Security* **3**(1), 1–7.
- Pötz, H., and Bleuze, P. (2012). *Urban Green-Blue Grids for Sustainable and Dynamic Cities*. Coop for Life.
- Ranger, N., Hallegatte, S., Bhattacharya, S., Bachu, M., Priya, S., Dhore, K., and Corfee-Morlot, J. (2010). An assessment of the potential impact of climate change on flood risk in Mumbai. *Climatic Change* **104**(1), 139–167. doi:10.1007/s10584-010-9979-2
- Rasul, G. (2012). Contribution of Himalayan ecosystems to water, energy and food security in South Asia: A nexus approach. ICIMOD. Accessed August 1, 2014: <http://lib.icimod.org/record/1898>
- Raudsepp-Hearne, C., Peterson, G. D., and Bennett, E. M. (2010). Ecosystem service bundles for analyzing tradeoffs in diverse landscapes. *PNAS* **107**, 5242–5247. doi:10.1073/pnas.0907284107.
- Register, R. (2006). *Ecocities: Rebuilding Cities in Balance with Nature*. New Society Press.
- Revi, A., Satterthwaite, D. E., Aragón-Durand, F., Corfee-Morlot, J., Kiunsi, B. R., Pelling Roberts, D. C., and Solecki, W. (2014). Urban areas. In *Climate Change 2014: Impacts, Adaptation and Vulnerability. IPCC Working Group II Contribution to AR5* (535–612). Cambridge University Press.
- Rogner, H. -H., Zhou, D., Bradley, R., Crabbé, P., Edenhofer, O. (Australia), B. H., and Yamaguchi, M. (2007). Introduction. In Metz, B., Davidson, O. R., Bosch, P. R., Dave, R., and Meyer, L. A. (eds.), *Climate Change 2007: Mitigation. Contribution of Working Group III to the Fourth*

- Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press.
- Rosenzweig, C., Solecki, W. D., Cox, J., Hodges, S., Parshall, L., Lynn, B., and Dunstan, F. (2009). Mitigating New York City's heat island: Integrating stakeholder perspectives and scientific evaluation. *Bulletin of the American Meteorological Society* **90**(9), 1297–1312. doi:10.1175/2009BAMS2308.1
- Rosenzweig, C., Solecki, W., DeGaetano, A., O'Grady, M., Hassol, S., and Grabhorn, P. (2011). *Responding to Climate Change in New York State: The ClimAID Integrated Assessment for Effective Climate Change Adaptation: Synthesis Report*. New York State Energy Research and Development Authority.
- Rosenzweig, C., Solecki, W. D., and Slosberg, R. B. (2006). *Mitigating New York City's Heat Island with Urban Forestry, Living Roofs, and Light Surfaces*. New York City Regional Heat Island Initiative.
- Rottle, N., and Yocom, K. (2011). *Basics Landscape Architecture 02: Ecological Design*. AVA Publishing.
- RUAF Foundation. (2014 March). Promoting urban agriculture as a climate change adaptation and disaster risk reduction strategy. *Urban Agriculture* 27.
- Salzman, J., Arnold, C. A., Garcia, R., Hirokawa, K. H., Jowers, K., LeJava, J., and Olander, L. P. (2014). The most important current research questions in urban ecosystem services. *Duke Environmental Law & Policy Forum* (1–47). Accessed June 20, 2015: http://scholarship.law.duke.edu/faculty_scholarship/3350
- Sandy Redevelopment Oversight Group. (2014). Equitable re-development of land acquired by New York City and New York State in Sandy-affected areas. Accessed May 27, 2015: <http://596acres.org/en/news/2013/12/19/what-our-new-mayor-should-do-redevelopment-in-sandy-affected-communities/>
- Santiago Fink, H. (2016). Human-nature for climate action: Nature-based solutions for urban sustainability, *Sustainability* **8**(3), 254. doi:10.3390/su8030254
- Satterthwaite, D., Huq, S., Pelling, M., and Reid, H. (2008). *Adapting to Climate Change in Urban Areas: The Possibilities and Constraints in Low- and Middle-Income Nations. Human Settlements Discussion Paper Series; Theme: Climate Change and Cities*. IIED.
- Satterthwaite, D., Huq, S., Pelling, M., Reid, H., and Romero, P. L. (2007 and 2008). *Adapting to Climate Change in Urban Areas: The Possibilities and Constraints in Low- and Middle-Income Nations. IIED working paper*. IIED.
- Schewenius, M., McPhearson, T., and Elmqvist, T. (2014). Opportunities for increasing resilience and sustainability of urban social-ecological systems: Insights from the URBES and the Cities Biodiversity Outlook Projects. *AMBIO*(Special Issue) **43**, 434–444. doi:10.1007/s13280-014-0505-z
- Scholz-Barth, K. (2001). Green roofs: Stormwater management from the top down. *Environmental Design and Construction*. January/February 2001, Posted on edcmag.com on 1/15/2001. Accessed December 17, 2014: <http://www.greenroofs.com/pdfs/archives-katrin.pdf>
- Searle, S. Y., Turnbull, M. H., Boelman, N. T., Schuster, W. S. R., Yakir, D., and Griffin, K. L. (2012). Urban environment of New York City promotes growth in northern red oak seedlings. *Tree Physiology* **32**, 389–400. doi:10.1093/treephys/tps027
- Secretariat of the Convention on Biological Diversity. (2012). *Cities and Biodiversity Outlook* (64 pages). Montreal.
- Seto, K. C., Parnell, S., and Elmqvist, T. (2013). A global outlook on urbanization. In Elmqvist, T., Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P. J., McDonald, R. I., Parnell, S., Schewenius, M., Sendstad, M., Seto, K. C., and Wilkinson, C. (eds.), *Urbanization, Biodiversity and Ecosystem Services: Challenges and Opportunities* (1–12). Springer. doi:10.1007/978-94-007-7088-1.
- Seto, K. C., Reenberg, A., Boone, C. G., Fragkias, M., Haase, D., Langanke, T., Marcotullio, P., Munroe, D. K., Olah, B., Simon, D. (2012). Urban land teleconnections and sustainability. *PNAS* **109**, 7687–7692. doi:10.1073/pnas.1117622109
- SMC Coastal Modeling System. (2015). Accessed February 9, 2016: <http://smc.ihcantabria.es/SMC25/en/>
- Solecki, W., and Marcotullio, P. J. (2013). Urbanization, biodiversity and ecosystem services: Challenges and opportunities. In Elmqvist, T., Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P. J., McDonald, R. I., and Wilkinson, C. (eds.), *Urbanization, Biodiversity and Ecosystem Services: Challenges and Opportunities* (485–504). Springer Netherlands. doi:10.1007/978-94-007-7088-1
- Steele, W., Mata, L., and Fünfgeld, H. (2015). Urban climate justice: Creating sustainable pathways for humans and other species. *Current Opinion in Environmental Sustainability* **14**, 121–126.
- Susca, T., Gaffin, S. R., and Dell'Osso, G. R. (2011). Positive effects of vegetation: Urban heat island and green roofs. *Environmental Pollution* **159**, 2119–2126. doi:10.1016/j.envpol.2011.03.007
- Suddick, E. C., and Davidson, E. A. (eds.) (2012). The role of nitrogen in climate change and the impacts of nitrogen-climate interactions on terrestrial and aquatic ecosystems, agriculture, and human health in the United States: A technical report submitted to the U.S. National Climate Assessment. Falmouth, MA: North American Nitrogen Center of the International Nitrogen Initiative (NANC-INI), Woods Hole Research Center.
- Sukhdev, P., Wittmer, H., Schröter-Schlaack, C., Nesshöver, C., Bishop, J., Ten, P., and Simmons, B. (2010). The Economics of Ecosystems and Biodiversity: Mainstreaming the economics of nature. A Synthesis of the approach, conclusions and recommendations of TEEB. Accessed April 12, 2016: http://www.urbanhabitats.org/v01n01/climatechange_full.html
- Sukopp, H., and Wurzel, A. (2003). The effects of climate change on the vegetation of Central European cities. *Urban Habitats* **1**(1), 66–86.
- Sutcliffe, S., and Court, J. (2005). *Evidence-Based Policymaking: What Is It? How Does It Work? What Relevance for Developing Countries?* Overseas Development Institute (ODI).
- Taha, H. (1997). Urban climates and heat islands: Albedo, evapotranspiration, and anthropogenic heat. *Energy and Buildings* **25**(2), 99–103. Accessed September 30, 2015: <http://www.sciencedirect.com/science/article/pii/S0378778896009991>
- Taylor, D. E. (2011). The evolution of environmental justice activism, research, and scholarship. *Environmental Practice* **13**(December), 280–301.
- The Economics of Environmentalism and Biodiversity (TEEB). (2010). *The Economics of Ecosystems and Biodiversity – Ecological and Economic Foundations*, ed. P. Kumar. Earthscan.
- The Economics of Environmentalism and Biodiversity (TEEB). (2011). TEEB Manual for cities: Ecosystem services in urban management. In UNEP and the European Union (ed.), *Ecosystem Services in Urban Management*. Accessed October 23, 2014: [http://www.teebweb.org/Portals/25/Documents/TEEB_Manual_for_Cities_Ecosystem_Services_for_Urban_management_FINAL_\(2011\).pdf](http://www.teebweb.org/Portals/25/Documents/TEEB_Manual_for_Cities_Ecosystem_Services_for_Urban_management_FINAL_(2011).pdf)
- Theurillat, J. -P., and Guisan, A. (2001). Potential impact of climate change on vegetation in the European Alps: A review. *Climatic Change* **50**(1–2), 77–109.
- Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C., and Williams, S. E. (2004). Extinction risk from climate change. *Nature* **427**(6970), 145–148. doi:10.1038/nature02121
- Troy, A. J., Grove, M., and O'Neil-Dunne, J. (2012). The relationship between tree canopy and crime rates across an urban-rural gradient in the great Baltimore region. *Landscaping and Urban Planning* **106**, 62–270.
- Tyler, S., and Moench, M. (2012). A framework for urban climate resilience. *Climate and Development* **4**(4), 311–326. Accessed April 10, 2016: <http://dx.doi.org/10.1080/17565529.2012.745389>
- Tzoulas, K., Korpela, K., Venn, S., Yli-Pelkonen, V., Kaźmierczak, A., Niemela, J., and James, P. (2007). Promoting ecosystem and human health in urban areas using Green Infrastructure: A literature review. *Landscape and Urban Planning* **81**(3), 167–178. doi:10.1016/j.landurbplan.2007.02.001
- UNEP. (2009). *The Role of Ecosystem in Climate Change Adaptation and Disaster Risk Reduction; Copenhagen Discussion Series*. UNEP.
- UNEP. (2011a). Supporting countries in meeting the climate challenge: UNEP's priorities for catalysing a green economy. Accessed May 27, 2014: www.unep.org/pdf/flagship_unep.pdf
- UNEP. (2011b). Towards a green economy: Pathways to sustainable development and poverty eradication – A synthesis for policy makers. Accessed October 13, 2015: www.unep.org/greeneconomy

- UNEP. (2012). UNEP Regional Seas Releases Report on Ecosystem Based Adaptation. Accessed August 28, 2014: <http://sdg.iisd.org/news/unep-regional-seas-releases-report-on-ecosystem-based-adaptation/?rdr=climate-l.iisd.org>
- UNFCCC Secretariat. (2011). Ecosystem-based approaches to adaptation: Compilation of information. Thirty-fifth session of the Subsidiary Body for Scientific and Technological Advice. Accessed August 13, 2015: <http://unfccc.int/resource/docs/2011/sbsta/eng/inf08.pdf>
- UN-Habitat. (2009). *Annual Report*.
- UN-Habitat. (2011). *Cities and Climate Change: Global Report on Human Settlements (2011)*. Earthscan.
- University of Cambridge and ICLEI. (2014). *Climate change: Implications for cities. Key findings from the Intergovernmental Panel on Climate Change Fifth Assessment Report*. Accessed December 16, 2015: <http://www.cisl.cam.ac.uk/Resources/Climate-and-Energy/Climate-Change-Implications-for-Cities>
- Urban, M. C., De Meester, L., Vellend, M., Stoks, R., and Vanoverbeke, J. (2012). A crucial step toward realism: responses to climate change from an evolving metacommunity perspective. *Evolutionary Applications* 5:154–167.
- Van Kerkhoff, L., and Lebel, L. (2006). Linking knowledge and action for sustainable development. *Annual Review of Environment and Resources* 31(1),445–477.
- Vignola, R., Locatelli, B., Martinez, C., and Imbach, P. (2009). Ecosystem-based adaptation to climate change: What role for policy-makers, society and scientists? *Mitigation and Adaptation Strategies for Global Change* 14, 691–696.
- Violin, C. R., Cada, P., Sudduth, E. B., Hassett, B. A., Penrose, D. L., and Bernhardt, E. S. (2011). Effects of urbanization and urban stream restoration on the physical and biological structure of stream ecosystems. *Ecological Applications* 21(6), 1932–1949. doi:10.1890/10-1551.1
- Von Döhren, P., and Haase, D. (2015). Ecosystem disservices research: A review of the state of the art with a focus on cities. *Ecological Indicators* 52(2015),490–497.
- Waldheim, C. (2007). Precedents for a North American landscape urbanism. In D. Almy and M. Benedikt (eds.), *Center 14 – On Landscape Urbanism* (292–303). Center for American Architecture and Design.
- Walker, M., Wilcox, B., and Wong, M. (2008). Waterborne zoonoses and changes in hydrologic response due to watershed development. In Fares, A., and ElKadi, A. (eds.), *Coastal Watershed Management* (349–367). WIT Press.
- Ward, P. J., Pauw, W. P., van Buuren, M. W., and Marfai, M. A. (2013). Governance of flood risk management in a time of climate change: The cases of Jakarta and Rotterdam. *Environmental Politics* 22, 518–536. doi:10.1080/09644016.2012.683155.
- Wardekker, J. A., de Jong, A., Knoop, J. M., and van der Sluijs, J. P. (2010). Operationalising a resilience approach to adapting an urban delta to uncertain climate changes. *Technological Forecasting and Social Change* 77, 987–998.
- While, A., and Whitehead, M. (2013). Cities: *Urbanisation and climate change*. *Urban Studies* 50, 1325–1331. doi:10.1177/0042098013480963
- Wilby, R. L., and Perry, G. L. W. (2006). Climate change, biodiversity and the urban environment: A critical review based on London, UK. *Progress in Physical Geography* 30(1), 73–98. doi:10.1191/0309133306pp470ra
- Wilkinson, C., Porter, L., and Colding, J. (2010). Metropolitan Planning and Resilience Thinking: A Practitioner's Perspective. *Critical Planning*. 17:2–20.
- Wilkinson, C., Sendstad, M., Parnell, S., and Schewenius, M. (2013). Urban governance of biodiversity and ecosystem services. In Elmqvist, T., Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P. J., McDonald, R. I., Parnell, S., Schewenius, M., et al. (eds.), *Urbanization, Biodiversity and Ecosystem Services – Challenges and Opportunities* (539–587). Springer Netherlands. doi:10.1007/978-94-007-7088-1.
- Williams, S. E., Shoo, L. P., Isaac, J. L., Hoffmann, A. A., and Langham, G. (2008). Towards an integrated framework for assessing the vulnerability of species to climate change. *PLoS Biology* 6, 2621–2626.
- Wisner, B., Blaikie, P., Cannon, T., and Davis, I. (2003). *At Risk: Natural hazards, people's vulnerability and disasters*, second edition. Routledge Taylor & Francis Group.
- Wolch, J. R., Byrne, J., and Newell, J. P. (2014). Urban green space, public health, and environmental justice: The challenge of making cities “just green enough.” *Landscape and Urban Planning* 125(1), 234–244.
- Woodall, C. W., Nowak, D. J., Likens, G. C., and Westfall, J. A. (2010). Assessing the potential for urban trees to facilitate forest tree migration in the eastern United States. USDA. Accessed November 8, 2015: http://www.nrs.fs.fed.us/pubs/jrnl/2010/nrs_2010_woodall_004.pdf
- Wortman, E., and Lovell, S. T. (2013). Environmental challenges threatening the growth of urban agriculture in the United States. *Journal of Environmental Quality*. doi:10.2134/jeq2013.01.0031
- Yang, J. (2009). Assessing the impact of climate change on urban tree species selection: A case study in Philadelphia. *Journal of Forestry* 107(7), 364–372.
- Yang, Z. (2013). *Eco-Cities: A Planning Guide*. Taylor and Francis Group.
- Yu, K., and Padua, M. (2005). *The Art of Survival: Recovering Landscape Architecture*. The Images Publishing Group Pty Ltd.
- Yu, K., and Li, D. (2006). The growth pattern of Taizhou City based on ecological infrastructure. In Yu, K., and Padua, M., *The Art of Survival (64–80)*. The Images Publishing Group.
- Zandersen, M., Jensen, A., Termansen, M., Buchholtz, G., Munter, B., Bruun, H. G., and Andersen, A. H. (2014). Ecosystem based approaches to Climate Adaptation – Urban Prospects and Barriers. Aarhus University, DCE – Danish Centre for Environment and Energy, 94 Scientific Report from DCE – Danish Centre for Environment and Energy No. 83. Accessed December 7, 2015: <http://dce2.au.dk/pub/SR83.pdf>
- Zavaleta, E. S., Shaw, M. R., Chiariello, N. R., Thomas, B. D., Cleland, E. E., Field, C. B., and Mooney, H. A. (2003) Grassland responses to three years of elevated temperature, CO₂, precipitation, and N deposition. *Ecological Monographs* 73, 585–604.
- Zeza, A., and Tasciotti, L. (2010). Urban agriculture, poverty and food security: Empirical evidence from a sample of developing countries. *Food Policy* 35, 265–273.
- Ziervogel, G., New, M., van Garderen, E.A., Midgley, G., Taylor, A., Hamann, R., Stuart-Hill, S., Myer, J., and Warburton, M. 2014. Climate change impacts and adaptation in South Africa. *WIREs Clim Change* 5:605–620.

Chapter 8 Case Study References

Case Study 8.1 Coastal Natural Protected Areas in Mediterranean Spain: The Ebro Delta and Emporda Wetlands

- Agencia Catalana de l'Aigua(ACA). (2014) Accessed April 17, 2015: http://acaweb.gencat.cat/aca/appmanager/aca/aca?_nfpb=true&_pageLabel=P51200158891423828204014
- Barnolas, M., and Llasat, M. C. (2007). A flood geodatabase and its climatological applications: The case of Catalonia for the last century. *Natural Hazards and Earth System Sciences* 7, 271–281.
- BirdLife International. (2014a). Important bird areas factsheet: Ebro Delta. Accessed November 8, 2015: <http://www.birdlife.org/datazone/sitefactsheet.php?id=1811>
- BirdLife International. (2014b). Important bird areas factsheet: Ampurdán marshes. Accessed December 7, 2015: <http://www.birdlife.org/datazone/sitefactsheet.php?id=1808>
- Bormann, H., Ahlhorn, F., and Klenke, T. (2012). Adaptation of water management to regional climate change in a coastal region – Hydrological change vs. community perception and strategies. *Journal of Hydrology* 454–455, 64–75.
- Candela, L., Fabregat, S., Josa, A., Suriol, J., Vigués, N., and Mas, J. (2007). Assessment of soil and groundwater impacts by treated urban wastewater reuse. A case study: Application in a golf course (Girona, Spain). *Science of the Total Environment* 374, 26–35.
- CIIRC. (2010). *Estat de la zona costanera a Catalunya*. CIIRC.

- Cote, M., Hurley, B., and Pratt, J. (2014). National Adaptation Planning: A Compendium of Stakeholder Workshops for USAID. Accessed June 26, 2015: http://pdf.usaid.gov/pdf_docs/PA00K2TB.pdf
- CREAF. (2013). Mapa de Coberturas del Suelo de Catalunya. Accessed September 30, 2014: <http://www.creaf.uab.es/mcsc/esp/index.htm>
- Day, J. W., Maltby, E., and Ibañez, C. (2006). River basin management and delta sustainability: A commentary on the Ebro delta and the Spanish national hydrological plan. *Ecological Engineering* **26**, 85–99.
- Eriksen, S., Aldunce, P., Bahinipati, C. S., D’Almeida Martins, R., Molefe, J. I., Nhemachena, C., O’Brien, K., Olorunfemi, F., Park, J., Sygna, L., and Ulsrud, K. (2011). When not every response to climate change is a good one: Identifying principles for sustainable adaptation. *Climate and Development* **3**, 7–20.
- Fatorić, S. (2010). *Vulnerability to the Effects of Climate Change and Adaptation: The Case of Ebro Delta*. Master’s thesis, Autonomous University of Barcelona.
- Fatorić, S. (2014). Vulnerability and Adaptation to Climate Change in the Mediterranean region: Climate out of balance in Aiguamolls de l’Empordà? Doctoral thesis, Autonomous University of Barcelona.
- Fatorić, S., and Chelleri, L. (2012). Vulnerability to the effects of climate change and adaptation: The case of Spanish Ebro Delta. *Ocean and Coastal Management* **60**, 1–10.
- Fatorić, S., and Morén-Alegret, R. (2013). Integrating local knowledge and perception for assessing vulnerability to climate change in economically dynamic coastal areas: The case of natural protected area Aiguamolls de l’Empordà, Spain. *Ocean and Coastal Management* **85**(A), 90–102.
- Fatorić, S., Morén-Alegret, R., and Kasimis, C. (2014). Exploring climate change effects in Euro-Mediterranean protected coastal wetlands: The cases of Aiguamolls de l’Empordà, Spain and Kotychi-Strofyliya, Greece. *International Journal of Sustainable Development & World Ecology* **21**(4), 346–360.
- Guillén, J., and Palanques, A. (1992). Sediment dynamics and hydrodynamics in the lower course of a river regulated by dams: The Ebro river. *Sedimentology* **39**, 567–579.
- IDESCAT. (2015). The municipality in figures. Accessed February 23, 2016: <http://www.idescat.cat/emex/?lang=en>
- Jiménez, J. A., Sánchez-Arcilla, A., Valdemoro, H. I., Gracia, V., and Nieto, F. (1997). Processes reshaping the Ebro delta. *Marine Geology* **144**, 59–79.
- Martín-Vide, J. P., Rodríguez-Máñez, E., Ferrer-Boix, C., Núñez-González, F., and Maruny-Vilalta, D. (2012). Estudio de la dinámica morfológica del río Fluvià. *Alcances y métodos frente a la escasez de datos. Tecnología y Ciencias del Agua* **III**(3), 115–133.
- Meteo Estartit. (2015). Accessed March 18, 2016: <http://www.historique-meteo.net/europe/espagne/estartit/2015/09/>
- Mozumder, P., Flugman, E., and Randhir, T. (2011). Adaptation behaviour in the face of global climate change: Survey responses from experts and decision makers serving the Florida Keys. *Ocean and Coastal Management* **54**(1), 37–44.
- Nelson, D. R., Adger, N., and Brown, K. (2007). Adaptation to environmental change: Contributions of a resilience framework. *The Annual Review of Environment and Resources* **32**, 395–419.
- Ninyerola, M., Pons, X., and Roure, J. M. (2004). Digital climatic atlas of Catalonia. Accessed July 24, 2014: http://www.opengis.uab.cat/acdc/english/en_cartografia.htm
- Observadora de l’Ebre. (2014). Accessed December 7, 2015: <http://www.observbre.es/en>
- Peel, M. C., Finlayson, B. L., and McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions* **4**(2), 462.
- Sánchez-Arcilla, A., Jiménez, J. A., Valdemoro, H. I., and Gracia, V. (2008). Implications of climatic change on Spanish Mediterranean low-lying coasts: The Ebro delta case. *Journal of Coastal Research* **24**(2), 306–316.
- SMC Coastal Modeling System. (2015). Accessed February 10, 2016: <http://smc.ihcantabria.es/SMC25/en>
- Smit, B., and Wandel, J. (2006). Adaptation, adaptive capacity and vulnerability. *Global Environmental Change* **16**, 282–292.
- World Bank. (2017). 2016 GNI per capita, Atlas method (current US\$). Accessed August 9, 2017: <http://data.worldbank.org/indicator/NY.GNP.PCAP.CD>
- Wetlands International. (1992). Delta del Ebro. Accessed March 11, 2015: http://sites.wetlands.org/reports/ris/3ES019en_FORMER_
- Wetlands International. (1995). Aiguamolls de l’Empordà. Accessed March 19, 2015: [http://sites.wetlands.org/reports/ris/3ES018en_FORMER_\(1995\).pdf](http://sites.wetlands.org/reports/ris/3ES018en_FORMER_(1995).pdf)

Case Study 8.2 New York’s Staten Island Bluebelt

- Agencia Catalana de l’Aigua (ACA). (2014) Accessed April 17, 2015: http://acaweb.gencat.cat/aca/appmanager/aca/aca?_nfpb=true&pageLabel=P51200158891423828204014
- Brauman, R. J., Gumb, D. F., and Duerkes, C. (2009). Designing for wildlife in the Blue Belt. *Clearwaters Winter* 2009, 41–43.
- Peel, M. C., Finlayson, B. L., and McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions* **4**(2), 462.
- Mayor’s Office, New York City. (2012). PlaNYC: A Stronger more resilient New York. Accessed August 16, 2015: <http://www.nyc.gov/html/planyc2030/html/home/home.shtml>
- New York City Department of Design and Construction, and the Design Trust for Public Space. (2005). High performance infrastructure guidelines. Accessed November 29, 2015: <http://www.nyc.gov/html/ddc/downloads/pdf/hpig.pdf>
- New York City Panel on Climate Change (NPCC). (2013). *New York City Panel on Climate Change Climate Risk Information 2013: Observations, Climate change projections, and maps*. City of New York.
- Ryan, G. C. (2006). *The return of the natives. The Urban Audubon* XXVII(4), 1,4.
- U.S. Census Bureau. (2010). Decennial census, summary file 1. Accessed November 17, 2014: <http://www.census.gov/population/metro/files/CBSA%20Report%20Chapter%203%20Data.xls>
- U.S. Census Bureau. (2016). Table 1. Annual Estimates of the Resident Population: April 1, 2010 to July 1, 2016 – Metropolitan Statistical Area; – 2016 Population Estimates. U.S. Census Bureau. Retrieved March 28, 2017. Accessed August 9, 2017: https://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=PEP_2016_PEPANNRES&prodType=table
- World Bank. (2017). 2016 GNI per capita, Atlas method (current US\$). Accessed August 9, 2017: <http://data.worldbank.org/indicator/NY.GNP.PCAP.CD>

Case Study 8.3 The Serra do Mar Project, Baixada Santista Metropolitan Region (BSMR), São Paulo State

- CDHU. (2012). Novos Bairros. Accessed March 9, 2014: http://www.agem.sp.gov.br/pdf/projeto_serra_do_mar_novos_bairros.pdf
- IAD, São Paulo. (2009). *Programa de Recuperação Socioambiental da Serra do Mar e do Sistema de Mosaicos da Mata Atlântica (Br-L1241). Estratégia Ambiental e Social do Programa. 05 de Outubro de (2009). Banco Interamericano de Desenvolvimento, Governo do Estado de São Paulo Brasil*. Accessed July 1, 2015: fflorestal.sp.gov.br/files/2012/02/easabidaserraadoamar.doc
- IBGE. (2015). *Instituto Brasileiro de Geografia e Estatística*. Accessed August 5, 2014: http://www.ibge.gov.br/home/estatistica/populacao/censo2010/sinopse/sinopse_tab_rm_zip.shtml
- IDB and São Paulo State Government. (2013). *Serra do Mar and the Atlantic Forest Mosaics System – A Social and Environmental Recovery Project. 1st Ed. Inter-American Development Bank (IDB), São Paulo State Government. São Paulo*. Accessed May 8, 2014: http://www.ambiente.sp.gov.br/serradomar/files/2014/03/INGL%C3%8AS_CARTILHA_Serra-do-Mar_Internet.pdf

- Peel, M. C., Finlayson, B. L., and McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions* 4(2), 462.
- São Paulo. (2013). Programa Serra do Mar e Mosaicos da Mata Atlântica (in Portuguese). Accessed May 5, 2014: http://www.agem.sp.gov.br/pdf/projeto_serra_do_mar_apresentacao.pdf
- World Bank. (2017). 2016 GNI per capita, Atlas method (current US\$). Accessed August 9, 2017: <http://data.worldbank.org/indicator/NY.GNP.PCAP.CD>

Case Study 8.4 Ecosystem-Based Climate Change Adaptation in the City of Cape Town

- Cilliers, S. S., and Siebert, S. J. (2012). Urban ecology in Cape Town: South African comparisons and reflections. *Ecology and Society* 17(3), 33. Accessed November 5, 2014: <http://dx.doi.org/10.5751/ES-05146-170333>
- City of Cape Town. (2015). City statistics. Accessed March 28, 2016: <https://www.capetown.gov.za/en/stats/Pages/CityStatistic.aspx>
- Laros, M. (2012). Cape Town, South Africa: An urban biodiversity network. ICLEI Case Study Series (part of the Local Sustainability). Kelleher, S., and Simpson, R., eds., ICLEI World Secretariat. Accessed August 15, 2014: http://www.iclei.org/fileadmin/PUBLICATIONS/Case_Studies/ICLEI_cs_138_Cape_Town.pdf
- Peel, M. C., Finlayson, B. L., and McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions* 4(2), 462.
- World Bank. (2017). 2016 GNI per capita, Atlas method (current US\$). Accessed August 9, 2017: <http://data.worldbank.org/indicator/NY.GNP.PCAP.CD>

Case Study 8.5 Jerusalem Gazelle Valley Park Conservation Program

- CIA World Factbook. (2015). Major Urban Areas – Population. Central Intelligence Agency. Accessed: August 4, 2017. <https://www.cia.gov/library/publications/the-world-factbook/fields/2219.html>
- Krasny, M. E. (2015). *Civic Ecology: Adaptation and Transformation from the Ground Up (Urban and Industrial Environments)*. MIT Press.
- Peel, M. C., Finlayson, B. L., and McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions* 4(2), 462.
- World Bank. (2017). 2016 GNI per capita, Atlas method (current US\$). Accessed August 9, 2017: <http://data.worldbank.org/indicator/NY.GNP.PCAP.CD>

Case Study 8.6 Medellín City: Transforming for Life

- Eveland, Jennifer. (2012). Medellín Transformed: From Murder Capital to Model City. Urban Redevelopment Authority. Accessed September 23, 2015: http://www.Leekuaneyewworldcityprize.com.sg/Features_Medellin_Transformed.htm
- Green City Index (GCI). (2010). Latin American Green City Index (2010). *Assessing the Environmental Performance of Latin America's Major Cities*. Siemens AG Corporate Communications and Government Affairs, Munich.

- Moreno, E. L. et al. (2013). *State of the World's Cities 2012/2013: Prosperity of Cities*. UN-Habitat, New York.
- Peel, M. C., Finlayson, B. L., and McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions* 4(2), 462.
- World Bank. (2017). 2016 GNI per capita, Atlas method (current US\$). Accessed August 9, 2017: <http://data.worldbank.org/indicator/NY.GNP.PCAP.CD>

Case Study 8.7 Singapore's Ecosystem-Based Adaptation

- Department of Statistics Singapore. (2015). Singapore in Figures 2015. Accessed January 18, 2016: [http://www.singstat.gov.sg/docs/default-source/default-document-library/publications/publications_and_papers/reference/sif\(2015\).pdf](http://www.singstat.gov.sg/docs/default-source/default-document-library/publications/publications_and_papers/reference/sif(2015).pdf)
- Department of Statistics Singapore. (2016) Yearbook of Statistics Singapore 2016. Accessed February 19, 2017: http://www.singstat.gov.sg/docs/default-source/default-document-library/publications/publications_and_papers/reference/yearbook_2016/yos2016.pdf
- Peel, M. C., Finlayson, B. L., and McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions* 4(2), 462.
- MEWR. (2014). Ministry of the Environment and Water Resources, The Singapore Green Plan 2012. Accessed July 9, 2015: [http://www.unccd\(2012\).org/content/documents/The%20Singapore%20Green%20Plan%20\(2012\).pdf](http://www.unccd(2012).org/content/documents/The%20Singapore%20Green%20Plan%20(2012).pdf)
- National Climate Change Secretariat Prime Minister's Office Republic of Singapore (NCCS). (2012). Climate change & Singapore: Challenges, opportunities, partnerships. Accessed July 24, 2014: [https://www.nccs.gov.sg/sites/nccs/files/NCCS-\(2012\).pdf](https://www.nccs.gov.sg/sites/nccs/files/NCCS-(2012).pdf)
- NEA. (2013). Climate Change, 2 September 2013. Accessed December 3, 2014: <http://www.nea.gov.sg/energy-waste/climate-change>
- World Bank. (2017). 2016 GNI per capita, Atlas method (current US\$). Accessed August 9, 2017: <http://data.worldbank.org/indicator/NY.GNP.PCAP.CD>

Case Study 8.8 Seattle's Thornton Creek Water Quality Channel

- Kaid Benfield's Blog. (2011). Outstanding urbanism, transit, and state-of-the-art green infrastructure, beautifully mixed In: SWITCHBOARD, Natural Resources Defense Council (NRDC) staff blog. Accessed October 2, 2014: http://switchboard.nrdc.org/blogs/kbenfield/outstanding_urbanism_and_state.html
- Peel, M. C., Finlayson, B. L., and McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions* 4(2), 462.
- Seattle Public Utilities. (2015). Thornton Creek water quality channel. Accessed March 14, 2016: <http://www.seattle.gov/util/MyServices/DrainageSewer/Projects/ThorntonCreekWaterQualityChannel/index.html>
- U.S. Census Bureau. (2010). Decennial census, summary file 1. Accessed September 23, 2014: <http://www.census.gov/population/metro/files/CBSA%20Report%20Chapter%203%20Data.xls>
- World Bank. (2017). 2016 GNI per capita, Atlas method (current US\$). Accessed August 9, 2017: <http://data.worldbank.org/indicator/NY.GNP.PCAP.CD>